

# Soils of the Bay of Plenty Volume 3

Eastern Bay of Plenty



Environment Bay of Plenty  
Environmental Publication 2010/11-3

5 Quay Street  
P O Box 364  
Whakatane  
NEW ZEALAND

ISSN: 1175 9372

*Working with our communities for a better environment  
E mahi ngatahi e pai ake ai te taiao*







# **Soils of the Bay of Plenty**

## **Volume 3: Eastern Bay of Plenty**

Environmental Publication 2010/11-3  
ISSN: 1175 9372

June 2010

Environment Bay of Plenty  
5 Quay Street  
PO Box 364  
Whakatane 3158  
NEW ZEALAND

---

Prepared by W. C. Rijkse and D. F. Guinto

---



# Preface

---

*Soil is a resource, a living, breathing entity that, if treated properly, will maintain itself.*

*It's our lifeline for survival. When it has finally been depleted, the human population will disappear.*

*Project your imagination into the soil below you next time you go into the garden. Think with compassion of the life that exists there. Think, the drama, the harvesting, and the work that carries on ceaselessly. Think about the meaning of being a steward for the earth.*

Marjorie Harris, *In the Garden* (1995)

For as long as I can remember I have been intrigued and fascinated by landscapes and soils of New Zealand, in particular of the Bay of Plenty where I spent a good deal of my career mapping soils. A landscape to me is a puzzle, a closed book, and to be able to open that book, to solve the puzzle by finding out what soils are in the landscape and what are the possibilities for good land use, is a joy we scientists call pedology – the science of soils. It is well developed in New Zealand, as befits a nation which traditionally earned so much of its wealth from the export of produce from the land. We are lucky in that respect that soils of the whole of the Bay of Plenty have now been mapped at a common scale of 1:50,000.

The general public can be excused for thinking that scientists are too pre-occupied with fine details of analysing soils for chemical and physical properties. We classify the soils, map their distribution and indicate their suitability for land use. Much of this knowledge was published by the Soil Bureau of the Department of Scientific and Industrial Research (DSIR) or, later, Landcare Research in reports, many of which are not available to the land user. This three-volume publication aims to make a major part of what we know about the soils of the Bay of Plenty available to farmers, foresters, horticulturists, farm consultants, and the general public.

We are lucky that Environment Bay of Plenty has the insight to finance the project of producing these soil books, and thus communicate soil knowledge to land users. I have enjoyed writing these books and call myself fortunate working with my colleague Dani Guinto who did more than his share producing these books. I hope that many of you will use this soil information to your advantage and getting the best from your land whilst realising your stewardship of the earth.

Wim Rijkse



## Acknowledgements

---

Many of the soil property data were derived from Landcare Research's S-map system, and we are grateful for that. Several photographs were taken from the soil survey of the Te Puke District, Bay of Plenty, New Zealand (unpublished) by W. E. Cotching. Acknowledgement goes to Simon Stokes of Environment Bay of Plenty for initiating this soil guide. The authors thank the many farmers and landowners in the Bay of Plenty for permission to enter their properties and to examine and photograph their soils.



# Contents

---

<b>Preface</b>	i
<b>Acknowledgements</b>	iii
<b>Part 1: Introduction</b>	1
<b>Part 2: The soil-forming environment</b>	3
2.1 Climate	3
2.2 Parent materials	3
2.2.1 Airfall tephra and flow tephra	4
2.2.2 Alluvium and colluvium	10
2.2.3 Organic materials	10
2.2.4 Wind-blown sand	10
2.3 Topography	10
2.4 Vegetation	11
2.5 Time	11
<b>Part 3: Soil landscapes</b>	13
<b>Part 4: Soil classification</b>	17
4.1 Soil orders in the Bay of Plenty	17
4.1.1 Allophanic soils	17
4.1.2 Anthropic soils	17
4.1.3 Brown soils	18
4.1.4 Gley soils	18
4.1.5 Organic soils	18
4.1.6 Podzols	18
4.1.7 Pumice soils	19
4.1.8 Raw soils	19
4.1.9 Recent soils	19

4.2	Soil horizons	20
<b>Part 5: Key soil management considerations</b>		21
5.1	Soil water retention and availability	21
5.2	Soil aeration and drainage	22
5.3	Irrigation	23
5.4	Leaching	24
5.5	Erosion	26
5.5.1	Surface erosion	27
5.5.2	Fluvial erosion	27
5.5.3	Mass movement erosion	27
5.6	Compaction	28
<b>Part 6: Guide to the soil series descriptions</b>		31
6.1	Soil series name	31
6.2	Overview	31
6.3	Physical properties	31
6.4	Chemical properties	35
6.5	Soil types/variations	35
6.5.1	Associated and similar soils	36
6.5.2	General land use suitability ratings	36
6.5.3	Management practices to improve suitability	36
6.5.4	Soil photos	36
<b>Part 7: Soils of Eastern Bay of Plenty</b>		37
<b>Part 8: Glossary</b>		233
<b>Part 9: References</b>		241
<b>Part 10: Index to Eastern Bay of Plenty soil series</b>		245

# Part 1: Introduction

---

The Bay of Plenty region covers approximately 21,836 km<sup>2</sup> (comprising 12,253 km<sup>2</sup> of land and 9,583 km<sup>2</sup> of coastal marine area) extending roughly from Katikati in the west to Cape Runaway in the east, and Rotorua District and parts of the Taupo District in the south.

Major landforms include:

- The low-lying Rangitaiki Plains, the plains near Cape Runaway on which river sediments have been laid down. Other flat coastal areas are located near Te Puke and Opotiki. Land use on the versatile soils on these landforms includes dairying, dry stock and horticulture.
- Much of the coastal land is on terrace-like flattish country on which thick layers of tephra occur. Land use includes dairying, dry stock and horticulture (kiwifruit, citrus, etc).
- The Rotorua Caldera was formed by large ignimbrite eruptions in the past. Land use on the tephric soils includes dry stock, dairying and horticulture, but much of the land is either residential or subdivided into small blocks.
- The Rotorua Caldera is flanked by the Mamaku Plateau where tephra overlies the ignimbrite which erupted out of the caldera. Land use consists of dry stock, some dairying and forestry.
- The Kaingaroa Plateau where tephra overlies ignimbrite. Land use is forestry.
- The Galatea Basin through which the Rangitaiki River flows consists chiefly of terrace-like surfaces covered by tephra. Land use consists of dairying and dry stock.
- Hill country forms much of the background of much of the above landforms.
- Steeplands (lands on slopes greater than 25 degrees) occur throughout the area covered with tephra, and dry stock and dairying are the main land uses. Large areas occur in the Urewera National Park and surrounding area. Much of this land is in indigenous forest or cut-over bush. Elsewhere, the steplands are used for dry stock farming.

Most previous soil mapping work in the Bay of Plenty was carried out by staff of the former Soil Bureau, a division of the Department of Scientific and Industrial Research. In later years some less published work was carried out by staff of Landcare Research. Publications of detailed soil survey work are listed in the References section.

Environment Bay of Plenty has combined all the major soil surveys and came up with a soil map of the whole region at a scale of 1:50,000 through Landcare Research's S-map system. This resource is available on the Internet and consists of a soil map and soil fact sheets (or soil reports) describing the physical and chemical properties of the soils. Access is through Environment Bay of Plenty's website (<http://www.envbop.govt.nz/Environment/Soil-Information.aspx>).

This publication is a companion resource to complement the soil information available online. In this publication, the soil information for the Bay of Plenty is provided in three volumes which describe the soils of Western, Central and Eastern Bay of Plenty. Figure 1.1 shows the three soil "sub-region" boundaries.




Figure 1.1 General soil map of the Bay of Plenty showing the soil orders and the sub-regional soil boundaries

These boundaries were drawn to minimise the occurrence of soil series common to two or three sub-regions (i.e. they are not based on district council boundaries). Where such overlaps occur (e.g. Matahina soil series occur in all three sub-regions), each soil series entry is repeated in each volume so that if readers are interested in one sub-region only, but their lands or properties of interest are at or close to the boundaries, they need not refer to the remaining volumes. When in doubt, however, it is advisable to consult all three volumes.

## Part 2: The soil-forming environment

---

The interactions among the principal factors of soil formation (parent material, climate, topography, vegetation and time) and soil-forming processes have given the soils of the Bay of Plenty their distinctive characters.

Parent materials range from thick layers of volcanic ash mantling the surface, to alluvium derived from greywacke, sandstone, mudstone and volcanic ash, to peat and wind-blown sand.

The climate of the Bay of Plenty varies from warm and moist in coastal areas to cool and moist in the uplands of Urewera National Park, the Mamaku Plateau and the Kaimai Range. It is probably the most important factor influencing present-day land use.

The influence of topography is somewhat subdued in a landscape mantled by tephra; however, strong dissection of hill country and steepland country influences the layers of tephra remaining on the slopes, and induces erosion and deposition of material on valley floors.

Vegetation has also played an important role in soil development. Changes in vegetation since the commencement of farming and commercial forestry have had considerable effects on properties such as soil stability.

Some of the principal environmental factors and their relationship to the soil pattern are discussed in greater detail below.

### 2.1 Climate

The Bay of Plenty is somewhat sheltered from the prevailing winds by the high country of the North Island. Consequently, the Bay of Plenty has a sunny climate with dry spells, but may have prolonged heavy rainfall periods.

Annual rainfall ranges from about 1,200 mm at the coast to over 2,000 mm inland at higher elevations, but decreases again in inland basins such as near Murupara.

Rainfall plays an important part in the development of soils. Broadly speaking, the higher the rainfall the stronger the leaching that takes place in the soil and, at annual rainfall over 1,800 mm, podzolisation processes are evident in the subsoil (redder subsoil).

Over 45% of the annual rainfall is recorded in the months from May to August. The driest period is from November to February. Seasonally, winter is generally the wettest and summer the driest part of the year (NZ Meteorological Service, 1973). Days with more than 1.0 mm rainfall range from around 110 a year at the coast to around 130 inland at Minginui Forest.

### 2.2 Parent materials

Basement rocks in southern parts of the Bay of Plenty consist chiefly of Urewera greywacke, argillite and basal massive, green volcanic sandstone of Jurassic and lower Cretaceous age with Pleistocene sandstone and siltstones south of Awakeri to Taneatua and west of Ruatoki, as well as south of Ohope as far as Waimana and east as far as the Raukokore River south of Cape Runaway. Past and present erosion resulted in generally shallow soils over angular, shattered greywacke, although tephra persists on stable ridges, crests and spurs. Little or no tephra remains on the steep slopes.

In western parts, ignimbrite and rhyolite form the main basement rocks, changing to andesite in the larger Katikati area.

Tertiary mudstones, siltstones, sandstones and gravels occur east of Waihou Bay. The northern part, east of Cape Runaway, is occupied by a large area of basalt.

Generally, these rocks are mantled by volcanic ash, and they are therefore not soil parent materials. However, especially on steep and very steep slopes, the kind of basement rock often determines the pattern and severity of erosion, and base rocks form an important component of the parent material of alluvium in areas such as the Rangitaiki Plains, the Opotiki flood plains and the flood plains near Cape Runaway.

The parent materials of the soils of the Bay of Plenty can be further divided into:

- Airfall tephra and flow tephra
- Alluvium and colluvium
- Peat
- Wind-blown sand

### 2.2.1 Airfall tephra and flow tephra

Volcanic eruptions occurred at different times from sources in Rotorua and Taupo Districts, and these eruptions were commonly violent, depositing coarse volcanic material called lapilli and blocks over the Bay of Plenty. Finer material or ash was usually deposited during the final stages of an eruption at greater distances away from the volcano. The general term used for all unconsolidated clastic volcanic material is tephra, and ash and lapilli refer to the grade or size only, as shown below:

fine ash	less than 0.25 mm
coarse ash	0.25 – 2.0 mm
lapilli	2 – 64 mm
blocks	more than 64 mm

Table 2.1 Volcanic ash showers in the Bay of Plenty

Ash shower	<sup>14</sup> C age in years before 1950 and volcanic centre	Occurrence and distribution	Characteristics
Tarawera Ash and Lapilli	64 Mt Tarawera	Western parts of Eastern Bay of Plenty and north of Kopuriki, thickness varies from 70 cm (Tarawera Forest) to 3 cm (near Whakatane).	Very dark greyish brown to dark grey sand and lapilli, lapilli varies in size from about 1 cm in Tarawera Forest to 1 mm near Whakatane.
Rotomahana Mud	64 Lake Rotomahana	Chiefly west and south of Mount Tarawera as far as Lake Rotorua and North towards the coast.	Greyish sandy loam to loamy sand layer at the surface.
Kaharoa Tephra	770 ± 20 Mt Tarawera	10 – 50 cm thick deposits in the northern part of southern and to the west of eastern Bay of Plenty.	Black sandy topsoils overlying pale yellow to white hard lapilli.
Taupo Tephra Taupo Ignimbrite	– Taupō	10 – 60 cm thick in the northern part of southern and to the west of eastern Bay of Plenty; the lower part of the horizon is rich in rhyolite.	Yellowish brown to pale yellow compact sand with few to many soft highly vesicular and fibrous lapilli and some dark grey rhyolite towards the base.
Taupō Lapilli	1850 ± 10 Taupō	Throughout the Bay of Plenty except towards Cape Runaway, varying in thickness from 60 – 70 cm in the southern parts to traces of lapilli in the north-east.	Pale yellow to strong brown uneven sized soft highly vesicular and fibrous lapilli.
Rotongaio Ash	– Taupō	Thin band in southern Bay of Plenty.	Dark grey to grey thin band which serves as a marker bed to indicate lower limit of Taupo lapilli. Not soil-forming.
Hatepe Lapilli	1,900 ± 60 Taupo	2 – 20 cm thick deposits in southern Bay of Plenty.	Light grey evenly sorted loose sand, not soil-forming but useful marker bed.
Waimihia Tephra Waimihia Ash	3,280 ± 20 Taupō	20 – 50 cm around Murupara and increasing in thickness south of Murupara.	Dark brown to dark yellowish brown, greasy gravelly sand to loamy sand.

Ash shower	<sup>14</sup> C age in years before 1950 and volcanic centre	Occurrence and distribution	Characteristics
Waimihia Lapilli	Taupō		Yellowish brown to dark yellowish brown, greasy gravelly sand, fine pumice gravel or sand, many fine lapilli. Strong brown colours at high altitudes.
Whakatane Tephra	4,830 ± 20 Ōkātina	50 – 60 cm thick through southern Bay of Plenty and western part of eastern Bay of Plenty, soil-forming in north-east and eastern parts of eastern Bay of Plenty.	Strong brown to yellowish brown slightly greasy loamy sand to sand with few to many pale yellow to yellowish brown medium lapilli.
Tuhua Tephra	6,130 ± 30 Mayor Island	30 – 40 cm below the topsoil, mostly north of Tauranga.	Yellowish brown sandy loam to silt loam.
Rotomā Tephra	8,530 ± 10 Ōkātina	Widespread in Bay of Plenty. About 200 cm at Matahina thinning out to 20 – 30 cm in Minginui Forest.	Yellowish brown greasy loamy sand, mostly in the lower subsoil.
Waiohau Tephra	11,850 ± 60 Ōkātina	Similar to Rotoma Tephra.	Yellowish brown greasy sandy loam to loamy sand in the subsoil.
Kawakawa Tephra	22,590 ± 230 Taupō	Various thickness but widespread in Northern Whakatane District.	Yellowish brown greasy sandy loam to loamy sand in the lower subsoil some andesitic ash (Tongariro) included in Southern Whakatane District.
Omataroa Tephra Mangaone Tephra Hauparu Tephra	28 220 ± 630 Ōkātina	Various thicknesses mainly south of the Rangitaiki Plains.	Thick layers of vesicular angular pale grey to pale yellow lapilli layers at several metres depth.
Rotoehu Tephra	ca. 50,000 Ōkātina	Various thicknesses throughout Bay of Plenty.	Distinctive white sand or lapilli at several metres depth.

Various descriptions of tephra occurring in the Bay of Plenty exist (Vucetich and Pullar 1969; Howorth 1975; Froggatt and Lowe 1990). The main ones are summarised in Table 2.1. Many of the listed tephra occur at depth, and only influence the soils where they are susceptible to erosion. For example, south of Ruatoki Valley, Rotoehu Ash overlies weathered greywacke and, where forest roads cut through the tephra layers, water accumulating in Rotoehu lapilli acts as a slide and, in saturated conditions, the whole tephra column slides off the greywacke. The most frequently occurring soil-forming tephra are Tarawera Tephra, Rotomahana Mud, Kaharoa Tephra, Taupo Tephra, Waimihia Tephra, Whakatane Tephra and Tuhua Tephra (See Figures 2.1 and 2.2).

Tarawera Tephra erupted from Mount Tarawera in 1886, and it occurs in topsoils from Galatea Basin towards the coast. It is thickest in the Tarawera Forest (70 cm) and it consists of black to dark greyish-brown angular basaltic ash and lapilli. It is considered to be significant where more than 7 cm thick.

Thickness and size of scoria can be critical on hilly slopes sown in pasture where coarse lapilli erode downslope and pastures fail to establish. The separation between the coarse textured Matahina series and the finer textured Manawahe series is based on such land use differences.

Rotomahana Mud erupted from the sides of Lake Rotomahana during the Tarawera eruption in 1886. The hydrothermally altered (“precooked” by thermal activity) material is quite fertile and supports excellent pastures.

Kaharoa Ash also erupted from Mount Tarawera (about 800 years ago) and it is widespread in central Bay of Plenty, except in southern parts. It consists of ash and white almost non-vesicular lapilli.

The Taupo Pumice Formation, hereafter called Taupo Pumice, represents the products of a series of violent eruptions which occurred shortly after each other in the Taupo area.

Taupo lapilli is the most widespread member of the Formation, and it occurs throughout the Bay of Plenty varying from at least one metre in the south to a few centimetres in the Rangitaiki Plains.

In central and southern areas it overlies Rotongaio Ash, a fine sandy hydrothermally altered mud, varying in thickness from 0.5 to 5 cm. This rests on Hatepe lapilli, a uniformly-graded lapilli (sago-like).

Taupo ignimbrite, formerly called Upper Taupo Pumice, overlies Taupo lapilli in central and southern Bay of Plenty. This material, called flow tephra, was deposited at the latter stages of the Taupo eruptions as a *nuée ardente*, or “glowing avalanche”, consisting of an incandescent mixture of ash and pumice moving rapidly like a gas cloud over the landscape. It left thick deposits of pumiceous sand, lapilli and blocks recognisable by poor sorting, compactness and the presence of many charred logs and pieces of charcoal. The lower part of the flow tephra contains much rhyolite, and it was formerly regarded as a separate layer called Rhyolite Block Member. Flow tephra is widespread and thick in large areas of Kāingaroa Forest south of its headquarters, and generally very thinly overlies rolling and hill country in southern Bay of Plenty.

The Waimihia Formation occurs in the subsoil of most soils in central and southern Bay of Plenty. Ash forms the upper part, and lapilli form the lower part. It is about 20 cm thick near Murupara, thickening to 100 to 200 cm in the south of the Bay of Plenty. It is a dark brown to dark yellowish brown greasy gravelly sand to loamy sand overlying yellowish brown to strong brown lapilli.




Figure 2.1 Tephra section north of Lake Rotoiti (Note: Numbers are years before present (1950)).




Figure 2.2 Present-day soil (Rotomahana silt loam), buried soil horizons and tephra layers at Brett Road, Rotorua (Source: Lowe 2006)

Whakatane Tephra is 50 to 60 cm thick in most areas and soil-forming in northern and north-eastern parts of Whakatane District where the younger tephra are thinning out. It consists of yellowish-brown to strong brown slightly greasy loamy sand to sand with few pale yellow to yellowish-brown medium lapilli.

Below Whakatane Tephra, various older tephra (and tephric loess) occur consisting of layers of yellowish-brown to pale yellow greasy loamy sand and sand, and layers of pumice lapilli occur with varying thickness. These tephra layers are generally not soil-forming on flat to rolling land, but on hilly and steep land, where the upper tephra have eroded off, they may occur closer to the surface. The tephra are listed in Table 1 with a general description of their occurrence and appearance in the field.

### 2.2.2 Alluvium and colluvium

Alluvium is widespread in areas such as the Rangitaiki Plains, the Opotiki area and generally on wide flood plains of the main rivers.

Much of the alluvium consists of rounded or sub-rounded pumice re-deposited from erosion products off hills. River alluvium frequently contains large amounts of material derived from sedimentary rocks. On the Rangitaiki Plains, much of the alluvium has been derived from Kaharoa Tephra and it often appears as a fine sandy or silty material (buff layer). It forms a compact layer of about 20 cm restricting root development and natural drainage. It overlies very thin to thin airfall Kaharoa Tephra. Alluvium in the Opotiki area is derived from tephra and greywacke, and near Cape Runaway, the alluvium is derived from greywacke and tephra with some mudstone and siltstone.

Colluvium is the product of erosion from hills accumulated on fans and valley floors. It is extensive in the eastern part of the Galatea Basin and on small fans throughout the Bay of Plenty. In the field it consists of much rounded and semi-rounded lapilli, and it looks often very similar to alluvium except that it tends to be coarser, e.g. contains more lapilli.

### 2.2.3 Organic materials

Peat occurs in many areas of the Rangitaiki Plains, on the Te Puke flats, and in the Opotiki area where it may occur in thick or thin layers alternating with pumice alluvium and/or airfall tephra. In some small areas, thin to very thin layers of diatomaceous earth occur in the subsoil.

### 2.2.4 Wind-blown sand

Wind-blown sand occurs in a belt along the coast and in local areas further inland. The dunes along the Rangitaiki Plains coast are covered or mixed with Tarawera Ash and Kaharoa Tephra. The tephra cover on the dunes further inland also includes Taupo Pumice and Whakatane Tephra.

## 2.3 Topography

Topography influences soil formation by modifying the climatic factor. By controlling the amount of runoff, topography influences the effectiveness of rainfall and the degree to which erosion removes the soil being formed. Similarly, the effectiveness of solar radiation varies with topography since the direction and steepness of slope determine the angle of incidence of the sun's rays. For example, in hill country, north-facing slopes generally receive more seasonal sunlight than south-facing slopes.

Low-lying areas that receive runoff retain most of the water from rainfall and are usually significantly wetter than neighbouring slopes. Variation in soil moisture regimes induced by differences in topographic positions is often the main reason for variation in soil properties over a landscape receiving a similar amount of rainfall per year. The effect of topography on the soil pattern at sub-regional level is discussed in the section on soil landscapes.

## 2.4 **Vegetation**

Evidence of pre-historic and pre-European vegetation is inferred from remnant forest pockets and buried wood and stumps. Vegetation patterns are closely related to the other soil-forming factors and soil type.

Dune lands are thought to have supported a forest of podocarp and kauri on the inland dunes and a mixture of pohutukawa and manuka on more coastal dunes. Well-drained terrace-like surfaces near the coast supported a mixed forest (rimu, tawa, kohekohe, kamahi and an occasional red beech or totara). Low-lying swampy areas supported kahikatea and pukatea swamp forest. Further inland, extensive podocarp forest (rata/tawa-rewarewa-mangeao-kamahi) changed to rimu-rata/tawa-kamahi forest where elevation and rainfall increased, such as on the Mamaku Plateau.

Today, much of the Bay of Plenty is sown into pasture or planted into subtropical fruit orchards along the coast, and pine forests further inland. Much of the indigenous forests remain, or are partly cut-over and allowed to re-grow, in the steep country of Eastern Bay of Plenty, and in the Urewera and Kaimai forests.

## 2.5 **Time**

Soil-forming factors and processes need time to produce soils. Relatively recently eroded steep slopes, and recent soils derived from alluvium or wind-blown sand, display a limited development of the subsoils. The soils tend to have low clay content, and sand particles dominate the soil matrix because of the limited time of weathering. Soil structures tend to be weakly developed and low levels of organic matter occur. On the other hand, soils derived from older parent materials such as tephra are more weathered, have greater clay content and deeper subsoil development compared with younger soils. Deposition of fresh sediments still continues today from erosion-deposition cycles on hilly slopes and along the main rivers and streams.

Although estimates of the rate of soil formation vary globally from 0.01 to 7.7 mm per year with an average of 0.1 mm per year, it is generally agreed that soil formation is a slow process (Morgan 1995). For example, most allophanic soils have taken between 10,000 and 20,000 years to form, and are clearly irreplaceable (Lowe and Palmer 2005). Thus, the importance of conserving topsoils cannot be overemphasised.



## Part 3: Soil landscapes

---

A soil landscape is a simplified representation of a sequence or pattern of soils in a landscape with respect to landforms or topographic positions. The soil landscapes are portrayed in two-dimensional (by distance and elevation) idealised cross sections of parts of the Bay of Plenty. They are more appropriately called schematic cross sections because in reality one or two soils may be missing in a given area and other soils may be more extensive. This publication uses the concept of soil series which may or may not contain two or more soil types. The schematic cross sections show selected soil types as they occur in soil landscapes. In the examples that follow, soils occurring within a soil landscape will vary from place to place with respect to slope, depth, texture, drainage, and other characteristics (e.g. depth of tephra layers). These differences in characteristics are important because they affect land use and soil management.

Figures 3.1 and 3.2 show the major soils of the Opotiki District. Figure 3.1 shows a cross section of the dunes from the fore dunes (Ohope), to soils with some topsoil (Papamoa) and the back swamps (Omarumutu). Behind the dunes, soils of the valley floors occur which frequently have poor drainage because of runoff from adjacent hill country (Opotiki hill soils).

Figure 3.2 lists the soils most frequently occurring on the Opotiki Plains. Beginning with a river channel in the west, it shows soils on low-lying river terraces that would flood frequently (Rangitaiki, Ruatoki, Kukumoa and Amokura) without stop banks. This leads to higher terraces with levees (Opouriao) and former back swamps (Otara, Apanui and Waioeka) which, without stop banks, would also flood more frequently. The next level up are tephra-covered terraces (Opotiki) with hilly slopes (Opotiki hill soils) to a narrow valley floor with imperfectly or poorly drained soils (Taho, Hanaia, Paerata and Iwiroa).

Figure 3.3 is a west–east section near Kawerau. It shows the correlation of soils with Tarawera Tephra in the upper layers. Soils with more than 20 cm Tarawera Tephra are Tarawera series, those with less than 20 cm coarse Tarawera Tephra are Matahina series and those with less than 20 cm fine Tarawera Tephra are Manawahe series. If soils have less than 7 cm Tarawera Tephra, which is difficult to see especially if ploughed, then they are mapped as Whakatane series.

Figure 3.4 shows a sequence of soils derived from Kaharoa Tephra. More than 30 cm of Kaharoa Tephra occurs in Pekepeke, Te Rere and Galatea series. At higher elevations, Ruakituri and Matawai series have less than 30 cm Kaharoa Tephra and are podzolised.




Figure 3.1 Schematic cross section of the major soils of Opotiki District showing the dune soils through to hill soils (Note: Diagram not drawn to scale).




Figure 3.2 Schematic cross section of the most frequently occurring soils in the Opotiki Plains (Note: Diagram not drawn to scale).




Figure 3.3 Schematic cross section of the soils of Whakatane District (Note: Diagram not drawn to scale).




Figure 3.4 Schematic cross section of the soils of southern Whakatane (Note: Diagram not drawn to scale).



## Part 4: Soil classification

---

The soils of the Bay of Plenty are discussed according to the New Zealand Soil Classification (Hewitt 1998a; 1998b). The classification has three categories of descending order: Order – Group – Subgroup. The series name can be regarded as a fourth level category in this classification. For example, Waitekauri series are Typic (subgroup) Orthic (group) Allophanic Soils (Order). In this publication, a classification to subgroup level is included in the overview. There are 15 soil orders in the New Zealand Soil Classification System. In the Bay of Plenty, nine soil orders are present (See Figure 1.1). These include: Allophanic Soils, Anthropic (or Man-made) Soils, Brown Soils, Gley Soils, Organic Soils, Podzols, Pumice Soils, Raw Soils and Recent Soils.

### 4.1 Soil orders in the Bay of Plenty

#### 4.1.1 Allophanic soils

Allophanic soils have properties that are strongly influenced by clay minerals that are poorly crystallised or amorphous (allophane, imogolite, ferrihydrite). They have weak soil strength and are sensitive with low bulk density. The soils are formed from layers of volcanic ash that are visible near the source (Rotorua area) and telescope together further away (East Coast area). They were called yellow-brown loams in previous soil classifications.

The soils typically have dark yellowish-brown grading to yellowish-brown sandy loam to silt loam subsoils with high levels of phosphate-fixing allophane in the clay fraction. Topsoils tend to be 18 cm or more deep with weakly developed structure and black to dark brown colours. The soils have a typically greasy feel when moistened and rubbed firmly between the fingers.

Allophanic soils are generally moderately to strongly leached with low levels of exchangeable calcium, potassium, magnesium and sodium. Reserves of magnesium and potassium are low to very low. Available phosphorus is naturally low with high phosphate retention.

Allophanic soils are friable to a great depth and do not have root-restricting layers. Along coastal Bay of Plenty, these are ideal soils for deep-rooting subtropical plants such as kiwifruit, provided they are sheltered from salt-laden winds. Other uses are pasture (dairying, dry stock), or forestry on steeper slopes. Cropping, such as maize, needs careful management to preserve topsoil structure.

#### 4.1.2 Anthropic soils

Anthropic soils or man-made soils are soils that have been altered by humans, including truncation of natural soils by earth-moving equipment, drastic mixing of natural soils so that their original character is lost, or by deposition of thick layers of organic or inorganic material.

Variation of areas thus affected is great and, in the Bay of Plenty, the worst cases are indicated on the soil maps but are not further described. Many soils were truncated in the 1980s in the kiwifruit growing areas, but such soils have not been mapped separately.

#### 4.1.3 **Brown soils**

Brown soils have secondary iron oxides evenly dispersed through the soil and give a yellowish-brown colour to the upper part of the subsoil. Base saturation values are usually moderate to very low. Poorly-drained soils are excluded from the order. Summer dryness and winter logging are uncommon.

Most of the brown soils occur in Eastern Bay of Plenty, where they support pasture (dry stock, dairying), forestry and cropping (maize). There are small areas in Western Bay of Plenty on steep slopes largely in indigenous forest.

#### 4.1.4 **Gley soils**

Gley soils are poorly or very poorly-drained in their natural state. Saturation occurs during prolonged periods, oxygen becomes limited, and reducing conditions occur. They were called gley soils or gleyed soils in previous soil classifications. Gley soils are essentially older wet mineral soils derived from a variety of parent materials including alluvium, colluvium and wind-blown coastal sands. They occur in valley floors, on the upper part of coastal terraces and coastal and river back swamps or former back swamps.

The soils have greyish-looking horizons, which have a lower boundary 90 cm or more from the mineral soil surface. It has low chroma (greyish) colours that occupy 50% or more of the matrix. Yellowish-brown or more reddish mottles are common. Dairying and dry stock are the main uses of gley soils. Maize is grown successfully on many gley soils.

#### 4.1.5 **Organic soils**

Organic soils have horizons that consist of organic material that, within 60 cm of the soil surface, has 30 cm or more peat accumulated in wet conditions. The soils were also called organic soils in previous soil classifications.

The soils occur in low-lying depressions or former back swamps and valley floors. They are poorly-drained with fluctuating water tables. High water tables reduce the rooting depth to about 20 cm in some areas. Many areas are drained with deep open drains resulting in some shrinkage of the peat. Land use includes dairying and dry stock, but pugging can be a problem during wet winters.

#### 4.1.6 **Podzols**

Podzols are strongly leached acid soils. They have a horizon of accumulation of aluminium occurring as complexes with organic matter and/or as short range minerals (typically as allophane/imogolite). The soils occur under high rainfall (generally exceeding 1,800 to 2,000 mm annually), mostly at higher elevations (exceeding 600 m), and they are usually associated with forest species which produce an acid litter, such as rimu and beech. Podzols are strongly leached and have low natural nutrient levels. The soils were called podzols or podzolised in previous soil classifications.

In the Bay of Plenty they are used for dry stock grazing and some dairying, forestry, and some cropping on the Mamaku Plateau. Large areas, such as the Urewera Range, are in indigenous forest.

#### 4.1.7 **Pumice soils**

Pumice soils are soils that are dominated by pumice or pumice sand high in volcanic glass. They have low clay contents and the clay fraction typically contains allophane. They were called yellow-brown pumice soils in previous soil classifications. Pumice soils occur in sandy or pumiceous tephra ranging from 700 to 3,500 years in age.

Clay contents are generally less than 10% and soil strength is weak or very weak. The soils are resistant to pugging and, like the Allophanic soils, have low to very low nutrient levels. The potential for erosion by water is high, especially when the surface vegetation and thin topsoil are removed. Summer droughts occur.

Pumice soils occur throughout Central Bay of Plenty in areas such as Rotorua, Kawerau and parts of the Rangitaiki Plains. Land use includes dry stock farming, dairying, forestry and fodder cropping.

#### 4.1.8 **Raw soils**

Raw soils lack distinctive topsoil development. They occur in environments where the development of topsoils is prevented by rockiness, active erosion, thermal activity, or deposition. They include beach sands or gravels.

The soils are mostly of medium fertility. They are well-drained with low water-holding capacity, and dry bulk densities above 1.4 tonnes per cubic metre. Main restrictions are frequent flooding, excessive drainage and extreme salinity. The soils are used for dry stock grazing or recreation.

#### 4.1.9 **Recent soils**

Recent soils show only incipient marks of soil-forming processes because of youthfulness, truncation of an older solum (the combined surface and subsurface horizons) or, less commonly, because the soil material is resistant to alteration. Soil formation has been sufficient to develop distinct topsoils. The concept of the order relates to weak soil development rather than the length of time of soil formation.

The main properties include weak soil development, generally high base saturation, gravel or rock not strongly altered, high potential rooting depth, good drainage, low phosphate retention, high fertility, and susceptibility to erosion and/or sedimentation.

Fluvial recent soils occur on slightly elevated river terraces. These are among the most versatile soils in the Bay of Plenty and land use includes dairying, dry stock and cropping (maize, fodder crops).

Recent soils also occur on hilly or steep slopes where surfaces are renewed after erosion. Land use on such slopes is mostly dry stock or forestry, the latter restricted because of shallow profiles overlying parent rock. Many areas are in indigenous forest.

## 4.2 Soil horizons

Soil horizon designations are used to indicate certain properties associated with a horizon. For example a topsoil is an A horizon. If that horizon has been ploughed or worked in any way, it becomes an Ap (p=plough) horizon. The following symbols have been used in the text:

A horizon	topsoil
Ap horizon	ploughed topsoil
Ah horizon	non-ploughed horizon
E horizon	bleached horizon immediately below the topsoil, occurs in podzols
Bw horizon	weathered horizon below the A horizon
Bh horizon	subsoil horizon with accumulation of organic matter and often iron and aluminium; occurs in podzols
Bg	gleyed or mottled (wet) horizon below the topsoil
C horizon	unconsolidated or weakly-consolidated mineral horizon that is little affected by soil-forming processes

## Part 5: Key soil management considerations

---

In this section, important management considerations for Bay of Plenty soils are discussed and generic soil management recommendations provided. Greater emphasis is given to managing the physical properties of soils for crop/animal production and environmental protection since they are more difficult to manipulate or change compared with chemical properties (e.g. nitrogen deficiency, high phosphate retention, acidity) which can be easily corrected by the application of fertilisers, lime, and other soil amendments. For plant growth an ideal soil has, as a rule of thumb, 50% solids, 25% water and 25% air by volume. From a physical standpoint soil management is about knowing one's soil and employing practices that optimise the relative proportions of air and water in the soil's pore spaces. The reader should consult more detailed soil management recommendations available in several excellent publications (e.g. Cornforth 1998; Shepherd 2009; Fertresearch nutrient management booklets: <http://www.fertresearch.org.nz/resource-centre/booklets>).

### 5.1 Soil water retention and availability

The retention and availability of water in the soil for plant use is largely determined by soil texture through its influence on the amount and size distribution of soil pores that retain water. Two useful measures of available water-holding capacity are **profile total available water** and **profile readily available water**. **Profile total available water** is the volume of water retained in the soil between field capacity and wilting point expressed as a depth (mm water per 100 cm soil depth). Field capacity may be defined as the water content that a soil reaches after it is saturated and allowed to drain until the drainage rate is negligible compared to evapotranspiration. It is considered the upper limit of the available water range (Figure 5.1). **Profile readily available water** is the volume of water expressed as a depth (mm water per 100 cm soil depth) considered extractable by plants with little effort and, consequently, little limitation to growth. It is the difference between soil field capacity and its water content when plant growth is inhibited by lack of soil water.

Available water-holding capacity is important for two reasons. First, it is a measure of the ability of the soil to sustain good growth and high yields of crops and pastures. Second, it is a measure of the capacity of the soil to store water via rainfall and irrigation (including effluent irrigation). The higher the available water-holding capacity, the more suitable the soil for any type of irrigation.




Figure 5.1 Soil water retention and depletion process illustrating the concepts of saturation, field capacity and wilting point. Available water-holding capacity is the amount of water retained between field capacity and wilting point (Source: University of Minnesota 2000).

Organic matter indirectly affects the water-holding capacity of soils by influencing soil structure and pore spaces. Organic matter stabilises soil structure and increases the volume and size of soil pores resulting in increase in water infiltration and water retention. Given the sandy nature of many soils of the Bay of Plenty, the importance of adding and/or retaining organic matter should be part of a sound soil management programme.


## 5.2 Soil aeration and drainage

Soil aeration reflects the ability of soil to allow exchange of air (particularly oxygen and carbon dioxide gases) between the atmosphere and plant roots. It represents the coarser pores in a soil, which provide the space into which plant roots grow and drain water from saturation to field capacity under the force of gravity. A soil is saturated when the total pore space is filled with water and the level of oxygen in the soil falls. Roots of most dryland plant species obtain oxygen used in aerobic respiration from surrounding pore space. This enables them to selectively absorb nutrients from the soil solution. When soil oxygen concentrations become limiting at saturation, the uptake by plants of major nutrients is inhibited, but manganese, iron and sodium may accumulate to toxic concentrations (Trought 1981). In waterlogged soil in winter time, plant growth may not be a great deal affected because oxygen requirements at that time of the year are small. Poor aeration and high moisture content also directly affect the occurrence and severity of some plant diseases, particularly in horticultural crops, e.g. phytophthora root rot (Cotching 1998).

Soil drainage refers to how much, and how quickly, water is removed from the soil. It also refers to the frequency and duration of periods when the soil is not wet. Drainage depends on three major factors: (1) input into the soil from rainfall, irrigation, seepage and runoff; (2) the flow of water through the soil (permeability); and (3) outlet from the soil or field drains to the collector drains, and from there to the sea (Griffiths 2004). Drainage is important because it affects both the oxygen supply and the temperature of the environment where plant roots and microorganisms thrive. Thus, wet soils will require some form of drainage if they are to be used productively for agriculture, since roots of non-aquatic plants do not normally survive and grow in saturated or waterlogged soils due to the lack of

oxygen. Plant growth on partially drained soils will be slow because wetter soils take longer to warm up so that growth does not start until later in the spring. Kiwifruit plants, for example, do not tolerate poor soil drainage. Animals grazing on wet land can compact the soil as they leave deep hoof marks that collect rainfall. Farm machinery may become bogged in wet paddocks and also compact the soil.

Most soil series of the Bay of Plenty have no or few problems with soil drainage as the soils are naturally well-drained. Exceptions include naturally poorly-drained gley and organic soils, which are generally located in coastal areas (Figure 5.2). The soil series descriptions, which form the main part of this publication, indicate whether drainage is a problem by listing the natural drainage class. In most areas in the Bay of Plenty open drains are used, because sandy or loamy subsoils are not suitable for mole drainage.



*Figure 5.2* Volume composition of a well-drained soil (Ōpōtiki sandy loam) and a poorly-drained soil (Apanui silt loam). Both soil types have almost equal pore space, but the volume occupied by air is much less in the poorly-drained soil because its pore space is largely occupied by water.

### 5.3 Irrigation

Despite the Bay of Plenty’s favourable annual rainfall for pasture and crop production, farmers are increasingly relying on irrigation to boost their productivity or avoid drought risks during the summer months. In particular, irrigation in kiwifruit has become increasingly common as more drought-prone soils are being used for this crop. Irrigation of effluent to soils is also becoming a common practice in dairy farms as a preferred method for treating effluent and for supplying water and nutrients to pasture.


Combined with climate information, knowledge of the water-holding capacity of soils is important (Section 5.1) to determine the correct amount and frequency of water application. Over-irrigation wastes water and energy and increases labour cost. It can also hasten the leaching of applied nutrients below the root zone (leading to contamination of water bodies), impede drainage, and reduce crop/pasture yields. On the other hand, under-irrigation stresses the plant and causes yield reductions. To avoid water stress, irrigation will need to be added to a level close to field capacity when half of the available water has been used by the plant. This is commonly known as the “refill point” or “trigger point”.

To avoid under- and over-irrigation, it is important to properly monitor soil moisture in the farm. Tensiometers, gypsum blocks, neutron probes, time domain reflectometry (TDR) and frequency domain reflectometry (FDR) sensors are the main instruments that can be used for monitoring soil moisture.

As water is becoming an expensive commodity nationwide, farmers should also investigate the use of precision irrigation technology (e.g. variable rate irrigation) which uses global positioning system (GPS) satellite guidance technology to take into account differences in soil and/or crop types that can be designated as separate management zones. Water (or fertiliser) application is tailored for each soil type (i.e. wetter soils receive less water than drier ones; fertile areas receive less fertiliser than infertile ones). Benefits include: savings in water, lower fuel consumption, application rates that are tailored for different soil and/or crop types, more efficient water/nutrient application, reduced nutrient runoff and leaching, less track maintenance, and evening out of inaccuracies in water distribution created where sprinkler nozzles are unable to apply the correct amount of water.

## 5.4 Leaching

Leaching is the process of removal of soluble materials (nutrients, metals and pesticides) in solution by water draining through the soil. The leaching potential of a soil depends on its texture, infiltration capacity, permeability, water-holding capacity, existing soil moisture content (representing a balance between rainfall or irrigation and the amount used by plants through evapotranspiration), concentration of soluble materials, and cation and anion exchange capacity. In addition, the movement of water down the soil profile can be non-uniform due to water flowing preferentially and more rapidly in cracks and other channels like worm holes, root holes, etc. (Figure 5.3). Such enhanced downward flow can carry a range of contaminants including nutrients, pesticides, trace metals, and pathogens in animal manure. Excessive leaching is detrimental since it leads to the loss of applied plant nutrients (e.g. nitrate-nitrogen and sulphate) resulting in greater fertiliser requirements/costs; water contamination when the leached nutrients and other unwanted substances reach the groundwater and eventually streams, rivers and lakes; and long-term acidification of soils due to the removal of basic cations like potassium, calcium and magnesium (Cornforth 1998). Strongly leached soils may occur anywhere, but are most common under high annual rainfall at higher elevations, as in the case of podzols.



*Figure 5.3 Applying water containing a blue dye stain to soil reveals uneven wetting of the soil profile due to water preferentially flowing in cracks and other channels. Nutrients carried by water, like nitrate, can potentially leach below the plant root zone and contaminate groundwater. (Photos from Carrick 2007)*

Nutrient leaching is very relevant to the Bay of Plenty because many of the soils have sandy textures. Nutrients in sandy soils with low organic matter content are easily leached because the soils are freely draining and have low nutrient retention capacities. On the other hand, soils containing appreciable clay and organic matter do not leach as much because a greater proportion of inorganic nutrients is adsorbed on the (mostly) negatively-charged exchange complex. Since texture is a basic soil property that is not easily changed, the addition of organic matter to sandy topsoils to increase their nutrient-holding capacity is very important. This applies largely to the nutrient cations potassium, calcium and magnesium which are held on the negative charges of organic matter. Nitrate anions are not held by the negative charges of organic matter, and can therefore still leach.

Leaching losses of nitrate can be reduced by judicious use of nitrogen fertilisers. The proper amount, method and timing of nitrogen fertiliser application are very important. Fertiliser applications should be scheduled to coincide with rapid plant growth to maximise nitrogen uptake. Where possible, applications should be split using smaller doses more often. It is essential to postpone fertiliser application if heavy rain is forecast or if the ground is too wet (near or above field capacity). Fertilisers should not be applied close to wetlands and streams. This will reduce potential for nutrients to be transported off-site during heavy rains. Fertiliser application should be reduced in areas where grazing animals congregate. The soils in these areas often contain sufficient nutrients for plant growth, so fertiliser application is often unnecessary.

## 5.5 Erosion

Accelerated soil erosion significantly affects the productivity and profitability of a farming enterprise both in the short and long term. It is therefore important to understand the processes, erosion types and soil conservation options available to the land manager. In New Zealand three broad categories of soil erosion are recognised, namely: surface erosion, fluvial erosion, and mass movement erosion. The categories are further subdivided into various erosion types (Lynn et al. 2009). Surface erosion involves the movement of a thin layer of soil particles or aggregates on the ground by the action of water, wind or gravity. Fluvial erosion involves the removal of soil material by running water flowing in channels. Mass movement erosion involves movement of soil and/or rock material downslope as a more or less coherent mass under the influence of gravity. Saturation of slope materials by water triggers mass movements.

Deposition, or sedimentation, may be regarded as a fourth erosion category as it is the endpoint of the soil erosion process, and can do as much damage as the removal of soil. Following is a discussion of the different types of erosion that commonly occur in the Bay of Plenty (Figure 5.4).




Figure 5.4 Common types of rainfall-induced soil erosion (Source: Auckland Regional Council 1999)

### 5.5.1 Surface erosion

**Sheet (or sheetwash):** Erosion in which thin layers of surface material are gradually removed more or less evenly from an extensive area of sloping land. It is caused by a combination of raindrop impact, detachment, and transport of soil particles by surface runoff. It is an insidious form of erosion as it removes topsoil and applied fertiliser.

**Scree (or scree creep):** The most common erosion type in the Bay of Plenty, whereby slow, gradual, more or less continuous, non-reversible deformation occurs, sustained by soil and rock materials under gravitational stresses. It occurs on many hill slopes and is recognisable by the presence of **terraces** on slopes. The steeper the slope, the more pronounced the terraces (see photos of Potikirua and Tawhia steepland soil landscapes in Volume 3).

**Wind:** Detachment, transport and deposition of loose materials by wind action. It occurs in coastal dune landscapes or along dry riverbeds, and may produce loess; however, loess is rare in the Bay of Plenty. Dry, volcanic ash soils are also very susceptible to this type of erosion. A good practice is avoiding cultivation during dry spells, which results in loss of organic matter.

### 5.5.2 Fluvial erosion

**Rill:** The formation of rills by water action. Rills are narrow (less than 30 cm wide) and shallow (less than 60 cm deep) channels that can be removed by tillage operations. However, when neglected, they can slowly develop into gullies.

**Gully:** The formation of gullies by water action. Gullies are channels which are wide (more than 30 cm) and deep (greater than 60 cm) enough to interfere with, and not be removed by, normal tillage operations. U-shaped gullies are common in the Bay of Plenty because of the relatively soft nature of tephra and alluvial deposits.

**Streambank (also called bank or channel erosion):** Erosion of soil material from the bank of a stream or watercourse caused by water flowing in stream and river channels, usually during periods of peak flow. Streambank erosion on recent alluvial soils can be very difficult to control as it is a natural process.

**Tunnel gully (also known as pipe/shaft erosion, under-runners or tomos):** Erosion by percolating water in a layer of subsoil resulting in tunnels or pipes which may collapse, producing gullies. Tunnel gullies may occur in thick subsoil pumice lapilli layers.

### 5.5.3 Mass movement erosion

**Slip:** This is a shallow, rapid sliding or flowing movement of soil material downslope, leaving a slip face and debris. The shearing takes place on a well-defined curved surface. It occurs during heavy rainfalls. Most slopes with a steepness of 15° can slip.

**Slump:** This tends to be deeper than a slip and involves downslope movement of large blocks of soil and rock materials. It involves rotational slide movements along curved planes of failure.

Recognising the different soil erosion types is important in planning and implementing appropriate soil conservation and management practices for sustaining agricultural productivity and protecting the environment. Depending on the location and soil type, different erosion types may be present in any given farm, and may require a combination of different soil conservation strategies. In general, preference is always given to biological or vegetative measures over mechanical measures, since they are less expensive and deal directly with reducing raindrop impact, increasing infiltration, reducing runoff volume, and decreasing water and wind velocities (Morgan 1995). Mechanical structures can be costly to install and maintain, and should be viewed as supplementing biological measures to control runoff flows. Surface soils of the Bay of Plenty are generally sandy and are regarded as erodible materials. The importance of maintaining adequate vegetative cover and the addition of organic matter to soils to reduce their susceptibility to erosion cannot be overemphasised.

Table 5.1 provides some general guidelines in combating the different types of erosion. Details of recommended soil conservation practices can be found in the Soil Conservation Technical Handbook published by the Ministry for the Environment (<http://www.mfe.govt.nz/publications/land/soil-conservation-handbook-jun01/soil-conserv-handbook-jun01.pdf>) and Environment Bay of Plenty's Land Management fact sheets (<http://www.envbop.govt.nz/Knowledge-Centre/Land-Management-factsheets.aspx>), or by contacting your local land management officer.

## 5.6 Compaction

An increase in the soil's bulk density or reduction in porosity is referred to as compaction. It is a process of packing the soil particles closer together causing a reduction in the volume of air. Compaction usually eliminates the largest air-filled pores first. Compaction can be caused by cultivation when the soil is wet, animal treading, and farm vehicular traffic. Driving on wet soils breaks down soil aggregates and compacts the soil. Excessive soil compaction restricts soil aeration, reduces plant growth and productivity, impedes drainage, reduces infiltration, and increases runoff generated during intense rains, leading to greater soil erosion losses.

Compaction can be minimised by waiting for the surface soil to dry out before driving on the soil, maintaining good soil structure and drainage, restricting the number of cultivation passes across the paddock, fitting dual wheels to reduce contact pressure and the risk of wheel slip, decreasing tyre pressures to reduce contact pressure, restricting heavy vehicles to the edge of the paddock (Cornforth 1998), practicing zero or minimum tillage, and the use of precision agriculture techniques.

There are few problems with compaction in the Bay of Plenty due to the sandy nature of many of the topsoils. However, compaction may occur in alluvial soils with clay loam topsoils that are used intensively, ploughed often, or when using heavy machinery with rubber tyres. The compaction is broken up by ripping. Compaction by animal treading under very wet soil conditions ("pugging") may also occur in soil types derived from Rotomahana Mud. Careful grazing management such as allowing the soil to dry below field capacity before grazing, limiting the number of hours animals graze on wet paddocks, the use of feed pads, animal shelters, etc. should be considered on such soil types.

*Table 5.1 Recommended soil conservation measures for various erosion types. (Adapted from Ngapo 2010)*

<b>Erosion type</b>	<b>Recommended soil conservation measures</b>
Sheet erosion	<ul style="list-style-type: none"> <li>• Maintain a good ground cover (vegetation or mulch) to protect soil surface from rain splash erosion.</li> <li>• Add organic matter to soils in the form of crop residues, composts, manures, cover crops, green manure crops, etc.</li> <li>• Practise zero or minimum tillage.</li> <li>• Avoid over-grazing; practice controlled grazing.</li> <li>• Site gates, fences, drinking troughs and other farm infrastructure carefully to avoid heavy concentration of stock on susceptible areas such as steep slopes.</li> <li>• Fence steep areas from flat areas.</li> <li>• Fence to slope aspect to allow for controlled grazing while avoiding over-grazing.</li> <li>• Use appropriate pasture species to suit the soil type.</li> <li>• Consider alternative land uses to grazing (e.g. protection forestry) of very steep slopes.</li> </ul>
Wind erosion	<ul style="list-style-type: none"> <li>• Avoid cultivation during dry spells.</li> <li>• Add organic matter to soils in the form of crop residues, composts, manures, cover crops, green manure crops, etc.</li> <li>• Practice zero or minimum tillage.</li> </ul>
Rill erosion	<ul style="list-style-type: none"> <li>• Stabilise the soil surface by maintaining a good ground cover of vegetation, mulch, and other materials.</li> <li>• Install runoff controls (e.g. graded banks, contour furrows and cut-offs) to reduce the velocity of overland flow. Ensure that runoff controls have a stable outlet.</li> <li>• Employ grass strips when cultivating.</li> </ul>
Gully erosion	<ul style="list-style-type: none"> <li>• Divert runoff away from the gully head.</li> <li>• Control runoff over the gully head using flumes, pipes and drop structures.</li> <li>• Reduce peak runoff rates.</li> <li>• Use a combination of the above methods (the most common method is to use a small detention bund, with adequate storm water detention, and low flow pipes to convey water over or around the gully head).</li> <li>• Employ planting of the gully head (for small gullies only).</li> <li>• Employ planting of stable points at critical locations.</li> <li>• Practise retirement of gullies from grazing in association with runoff control.</li> <li>• Contour ground to “smooth out” small gullies on low terraces (in combination with runoff control and surface vegetation).</li> </ul>

Erosion type	Recommended soil conservation measures
Streambank erosion	<ul style="list-style-type: none"> <li>• Avoid grazing of livestock on streambanks.</li> <li>• Fence and plant streams with careful thought on the type and location of fences and what planting should be done (grass buffers and native species are preferred).</li> </ul>
Tunnel gully erosion	<ul style="list-style-type: none"> <li>• Plant a willow pole in the tunnel hole.</li> <li>• Collapse in with a shovel or digger to remove the risk of lamb or sheep losses.</li> <li>• Bulldoze in or fill the hole with coarse material to filter out any fines.</li> <li>• Fence the tunnel gully to ensure stock do not fall in.</li> </ul>
Slips and slumps	<ul style="list-style-type: none"> <li>• Plant soil-conserving trees (e.g. poplars; willows; native trees like manuka, kanuka).</li> </ul>

## Part 6: Guide to the soil series descriptions

---

The soils are listed alphabetically according to soil series. The following is a guide to understanding the soil information contained in the soil series descriptions. Soil-related terms are also described in the Glossary.

### 6.1 Soil series name

This is the name of the soil series and its corresponding soil map symbol or code. A soil series is a group of soils that have similar profile characteristics except for differences in texture of the surface layer. Traditionally, a soil series is named after the place where the soil was first observed and described (e.g. Matahina series).

### 6.2 Overview

The overview provides general information on the occurrence and distribution of the soil series, parent materials, physiographic position and slope, colour, profile texture, soil classification (up to subgroup level), vegetation, and/or land use. Where relevant, occurrence of soils under varying rainfall amount is also mentioned (e.g. in the case of Podzols).

It should be noted that soils are rarely homogenous. This is particularly the case with steepland soils which have highly variable characteristics. Thus, if a steepland soil is classified as an allophanic soil, this is a generalisation. In reality, the soil is predominantly allophanic but, when it occurs on steep, eroded slopes, it can also be classified as a recent soil.

### 6.3 Physical properties

**Texture:** Texture is the relative proportions of the primary particles in the soil, namely sand (2.00-0.06 mm), silt (0.06-0.002 mm) and clay (<0.002 mm). Particles with diameters less than 2 mm are referred to as the fine earth fraction, while those with diameters greater than 2 mm are called coarse fragments. Sand particles feel gritty, and are large enough to be seen by the naked eye. Silt particles are smaller, and feel smooth, like flour. Clays are much smaller, feel sticky, and can be formed into ribbons and wires when wet. Clay particles are flat, and can be seen only when viewed under a high-powered microscope. Because sand and silt are relatively large, they possess only a small surface area and contribute little to the chemical behaviour of the soil. In contrast, clays are very small, possess a large surface area, carry electrical charges, and are much more chemically active than sand and silt.

Every soil contains a mixture of sand, silt and clay, and this is expressed as a textural class name such as sandy loam, silt loam, clay, etc. A soil that contains a balanced mixture of sand, silt and clay is called a loam. If rock or stone fragments are present in significant quantity, then a coarse fragment modifier, such as gravelly sandy loam, is used. The soil textural triangle (Figure 6.1) shows all the textural class names that result in various combinations of sand, silt and clay. A more generalised, or simplified, textural triangle is used to group the textural classes into sandy, silty, loamy and clayey.

The texture given here actually refers to the simplified textural groupings. Skeletal soils (those with horizons containing 35% or more gravel by volume) and peats are also included as textural groups. A soil can have two or more contrasting textures with depth and is described as layered (e.g. loam over sand, loam over clay, sand over skeletal). In the Bay of Plenty many soils are formed from volcanic deposits, and the 'gravel' consists of pieces of pumice called lapilli.

Texture is a basic property of the soil that is not easily changed. It affects other soil properties, such as water availability, permeability, drainage, and aeration. It also influences nutrient retention, the development of soil structure, and the ease of soil cultivation. Texture influences the balance between water-filled pores and air-filled pores, creating different soil environments for root growth and the activity of microorganisms. Medium-textured soils, such as loam and silt loam, have a range of pore sizes that allows water to flow through the smaller pores and exchange air in the larger pores. Thus, they provide favourable environments for root growth, store large amounts of water for plant use, and have good nutrient-supplying power. Sandy soils have more large pores and fewer small pores. They have good aeration, but store much less water for plant use, and are considered droughty soils. In heavily-fertilised sandy soils rapid water movement increases the risk of groundwater pollution through leaching of excess nutrients such as nitrate.




Figure 6.1 Soil textural triangles showing the textural classes (left) and simplified textural groups (right)

**Topsoil clay content:** This refers to the range of clay content of the topsoil expressed as a percentage. In general, soils of the Bay of Plenty do not contain appreciable amounts of clay. Clay particles are very small with a high surface area per unit weight, giving them a high capacity to adsorb water and other substances. They also carry electrical charges (mostly negative), and so they are the most chemically active part of the soil. Clay is responsible for the swelling and shrinkage of the soil, and makes moist soil sticky. Positively-charged plant nutrient elements (cations), like ammonium, calcium, magnesium and potassium, are adsorbed on the negatively charged surfaces of clays, which helps retain these nutrients in the soil for plant use.

In addition to the amount of clay present in the soil, the type of minerals present also influences soil behaviour. Common clay minerals in Bay of Plenty soils include halloysite or kaolinite, montmorillonite, and allophane. A soil will generally have varying proportions of minerals in the clay fraction. However, the clay fraction of some soils derived from volcanic ash is dominated by allophane – a non-crystalline mineral which imparts good soil structure and high water-holding capacity, but is capable of fixing large amounts of phosphorus (P) rendering it unavailable for plant use.

**Potential rooting depth:** This refers to the soil depth (in cm) to the top of a barrier (within 1 m of the soil surface) that limits root extension.

**Rooting barrier:** This is the type of barrier that limits root extension. Root penetration into soil may stop due to a physical barrier (e.g. when it encounters a compacted layer, poor aeration, low moisture content), a chemical barrier (e.g. aluminium toxicity in the subsoil), or a combination of these.

**Drainage class:** The drainage class indicates how long a soil or part of a soil is saturated with water, and how quickly it can drain excess water. Some soils have mottles, which are spots of grey or brown colour different from the main colour of the soil. Mottles indicate the height of fluctuating water tables. Grey mottles are indicative of waterlogging and reduction of iron compounds in the soil. Drainage of the soil is important so that the pore spaces are able to be filled with oxygen for plant roots to access. Simplified descriptions of soil drainage classes are shown in Table 6.1. A more detailed definition of drainage classes is given in Milne et al. (1995).

Table 6.1 Simplified soil drainage classes

Class	Description
Very poorly-drained	Organic-enriched topsoil (peaty) with grey subsoil
Poorly-drained	Grey layer begins just below the topsoil
Imperfectly drained	Grey layer at 40 – 90 cm depth or mottling within 30 cm of the surface
Moderately well-drained	Grey layer at 60 – 90 cm depth or mottling between 30 and 90 cm
Well-drained	No grey layer or mottling within 80-90 cm depth

**Permeability:** This is a measure of the rate at which water can flow through the soil. Permeability is dependent on the amount, size, shape and interconnectedness of soil pores which are influenced by soil texture, soil structure, and soil organic matter. Sandy soils have larger pores and more rapid permeability than clayey soils. Good soil structure promotes high permeability by providing stable aggregates consisting of small pores within the aggregates and large pores between them. Organic matter increases permeability through its binding action on soil aggregates. A soil's overall permeability is usually based on the horizon with the slowest permeability class and the depth at which this layer occurs. Permeability is important for ease of drainage, risk of waterlogging, effluent absorption potential, leaching, and water loss. Permeability classes include: slow (less than 4 mm/hr), moderate (4 – 72 mm/hr) and fast (greater than 72 mm/hr).

**Profile total available water (0-100 cm):** This is the amount of water (in mm) that can be extracted between field capacity (-10 kPa suction) and permanent wilting point (-1500 kPa suction) to a depth of 1 m. Profile total available water is important for droughtiness and overall water availability. Classes are shown in Table 6.2.

Table 6.2 Profile total available water classes

Class	Range (mm)
Low	30 – 60
Low to moderate	60 – 90
Moderate	90 – 120
Moderate to high	120 – 150
High	150 – 250
Very high	250 – 350

**Profile readily available water (0-100 cm):** This refers to the amount of water (in mm) held in a soil that can be easily extracted by plant roots within the potential rooting depth (i.e. between field capacity (-10 kPa suction) and -100 kPa suction). Classes are shown in Table 6.3.

**Topsoil and subsoil bulk density:** This is the dry mass of the fine earth fraction (<2 mm) divided by the total soil volume and is expressed in grams per cubic centimetre ( $\text{g}/\text{cm}^3$ ), or tonnes per cubic metre ( $\text{t}/\text{m}^3$ ). The total or bulk volume consists of the volume of soil solids and the volume of soil pores. Thus, bulk density is a measure of the degree of soil compaction since it includes the volume of pores. Compacting a soil results in a lower volume occupied by pore spaces resulting in higher bulk density. Therefore the higher the bulk density, the lower the porosity, and the slower the drainage.

Table 6.3 Profile readily available water classes

Class	Range (mm)
Very low	<25
Low	25 – 50
Moderate	50 – 75
Moderate to high	75 – 100
High	100 – 150
Very high	150 – 250

Bulk density affects available water content and air capacity of soils and is an indicator of the ease of root penetration. Dry bulk density tends to be higher in soils with higher clay content. Dry bulk densities greater than  $1.6 \text{ g}/\text{cm}^3$  are likely to be associated with high strength, and may represent an impediment to root penetration. Those lower than  $0.4 \text{ g}/\text{cm}^3$  are probably associated with material of recent volcanic origin, and are likely to cause engineering problems. Dry bulk densities of less than  $0.2 \text{ g}/\text{cm}^3$  indicate that considerable shrinkage will occur when these materials are drained, as in the case of peat.

## 6.4 Chemical properties

**Topsoil organic matter:** This is the amount of organic matter in the topsoil expressed as a percentage. The typical dark colour of many surface soils is due to organic matter. Organic matter can hold up to 20 times its weight in water, and so improves the water-holding capacity of soils, particularly drought-prone sandy soils. Organic matter cements soil particles into structural units called aggregates, which stabilise soil structure, improve aeration, and increase permeability. Like clay particles, organic matter possesses a high surface area and lots of negative charges. These negative charges attract positively-charged ions (cations), such as calcium, magnesium, potassium, etc., which would otherwise leach in the soil profile. Organic matter is also an important source of plant nutrients. When it decomposes it releases the major plant nutrient elements nitrogen, phosphorus and sulphur. Organic matter exhibits chemical buffering, which helps the soil resist rapid changes in pH. Finally, organic matter plays a crucial role in sequestering carbon from the atmosphere. Because of its many production and environmental benefits, the maintenance of a high level of soil organic matter is essential in any sustainable soil management programme.

**Topsoil pretention:** Expressed as a percentage, it is a measure of the ability of the soil to remove phosphorus (phosphate) from the soil solution, rendering it unavailable to plants. In acidic soils phosphate is retained by reactive aluminium and iron minerals. In alkaline or calcareous soils phosphate is precipitated as calcium phosphate compounds. In allophanic soils, phosphate is retained by allophanic minerals. High P retention indicates that plants will give a lower response to the same amount of phosphate fertiliser than on a soil with low P retention. This is often the case with allophanic and some pumice soils. Phosphate retention values influence phosphate fertiliser requirements and soil structural stability. P retention classes are shown in Table 6.4.

Table 6.4 *Topsoil P retention classes*

Class	Range (%)
Very low	<10
Low	10 – 30
Medium	30 – 60
High	60 – 90
Very high	>90

**Available P, Ca, Mg and K:** This provides a general indication of the levels of available phosphorus, calcium, magnesium and potassium in the soil.

## 6.5 Soil types/variations

This lists the soil type(s)/variations present within the soil series. The soil type is a subdivision within a soil series to distinguish soils differing in surface texture only (e.g. Taupo sand, Taupo sandy loam, etc.). Phases of soil series which are based on features that affect the use and management of the soil, such as slope, depth, stoniness, etc., are also included. For example, slope steepness is an important factor determining land use, so hill soils (16 – 25 degrees) and/or steepland soils (greater than 25 degrees) are distinguished in some soil series.

### 6.5.1 **Associated and similar soils**

This indicates the soils that are geographically associated with the soil series (e.g. soils occurring on similar topographic positions).

### 6.5.2 **General land use suitability ratings**

General land use suitability ratings are provided for arable, horticulture, intensive pasture, and forestry land uses. Ratings for each land use include: not suitable, low, moderate, and high. The *management considerations* portion addresses the limitations posed by a particular soil series (e.g. steep slopes, low fertility, etc.).

### 6.5.3 **Management practices to improve suitability**

This provides a generic guide to soil management practices to overcome soil limitations and improve suitability of the soil for the various land uses considered.

### 6.5.4 **Soil photos**

Soil profile photos of most soil series are included. For other soil series on steep or waterlogged areas, photos of soil landscapes are shown instead.

## Part 7: Soils of Eastern Bay of Plenty

---

**Soil Series Name:** Amokura (Am)

### Overview

Amokura soil series occur in the Opotiki area in small depressions (oxbows) along the active flood plains of the main rivers (Waioeka, Otara, Waiaua and Waiotahi Rivers). Soil profiles, formed from alluvium, have thin mottled silt loam topsoils overlying olive grey or very dark greyish brown mottled silt loam subsoils. They are classified as **Acidic Recent Gley Soils**. The soils generally support rough pastures, some improved pasture and small areas in maize.

### Physical properties

**Texture:** Silt loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** Unlimited

**Rooting barrier:** Water table

**Drainage class:** Poorly drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** High (178 mm)

**Profile readily available water (0 – 100 cm):** High (100 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.0 – 13.8%

**Topsoil P retention:** Medium (35%)

**Available P, Ca, Mg and K:** Low to medium

### Soil types/variations

Topsoil textures range from **silt loam** to **fine sandy loam** and some subsoils are silty clay loam.

### *Associated and similar soils*

**Rangitaiki series** occur on the same river terrace but they are well drained. **Kukumoa series** are imperfectly drained with deeper topsoils.

## **General land use suitability ratings**

### **Amokura silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Poor drainage, flooding risk.
Horticulture	Not suitable	Poor drainage, flooding risk.
Intensive pasture	Low to moderate	Poor drainage, risk of pugging.
Forestry	Not suitable	Poor drainage, flooding risk.

### **Management practices to improve suitability**

- Employ surface drainage (but areas are often too small to economically drain).



**Amokura silt loam**

## Soil Series Name: Apanui (Ap)

### Overview

Apanui soil series are derived from alluvium of former back swamps of main river terraces in the Opotiki area. Soils have deep (20 to 22 cm) silt loam topsoils overlying mottled grey clay loams and clays. A 10 cm thick layer of fine Taupo Tephra occurs at 50 cm depth. The soils are classified as **Typic Orthic Gley Soils**. They are used for intensive pastoral use (dairying), some dry stock, cropping (maize), and horticulture (cabbages, cauliflower).

### Physical properties

**Texture:** Loam over clay

**Topsoil clay content:** 38 – 42%

**Potential rooting depth:** Unlimited

**Rooting barrier:** None

**Drainage class:** Poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Moderate (120 mm)

**Profile readily available water (0 – 100 cm):** Moderate (71 mm)

**Topsoil bulk density:** 0.87 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.01 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.0 – 13.8%

**Topsoil P retention:** Medium (35%)

**Available P, Ca, Mg and K:** P high to medium; Ca low; high reserves of Mg and K

### Soil types/variations

Topsoil textures vary from silt loam to clay loam.

#### ***Associated and similar soils***

**Waioeka series** have layers of diatomaceous earth. **Otara series** are imperfectly drained and occur on the margins of former back swamps. They generally have browner (rather than grey) colours in the subsoil.

## General land use suitability ratings

### Apanui silt loam

Land use	Suitability rating	Management considerations
Arable	Moderate	Poor drainage, locally heavier textured topsoil (clay loam).
Horticulture	Moderate	Poor drainage, locally heavier textured topsoil (clay loam).
Intensive pasture	Moderate to high	Poor drainage.
Forestry	Not suitable	Poor natural drainage.

### Management practices to improve suitability

- Drainage: Employ surface drainage or mole drainage.
- Plant shallow-rooting crops for horticulture (cabbages, cauliflower).



**Apanui silt loam** (Note subsoil mottling and Taupō Pumice layer at 50 cm depth)

## Soil Series Name: Awaiti (Ati)

### Overview

Awaiti soil series occur on the flat, higher parts of the former flood plain of the Tarawera River on the Rangitaiki Plains. They are derived from thin and very thin Tarawera Tephra on fine pumiceous alluvium derived from Kaharoa Tephra. Soil profiles have black topsoil (with fine Tarawera lapilli) over dark grey silt loam with a sharp boundary to pale brown sand. The soils are classified as **Buried-pumice Tephric Recent Soils**. Present land use is dairying with fodder cropping.

### Physical properties

**Texture:** Sandy loam over sand and silt loam

**Topsoil clay content:** 10 – 18%

**Potential rooting depth:** Unlimited

**Rooting barrier:** Fluctuating water table

**Drainage class:** Well drained to moderately well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** moderate (102 mm)

**Profile readily available water (0 – 100 cm):** moderate (67 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.38 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 - 12%

**Topsoil P retention:** Medium (32%)

**Available P, Ca, Mg and K:** Low to medium

### Soil types/variations

**Awaiti sand** is the main soil type. Yellowish brown mottles may occur in the subsoil. Inclusions of **Kawerau series** and **Omeheu series** occur near the boundaries of the soils.

#### ***Associated and similar soils***

**Te Teko series** have more organic matter in the topsoil and less sandy in the subsoil. **Kawerau series** have coarse sandy subsoils and generally coarser Tarawera lapilli in the topsoil. **Omeheu series** have similar parent material but poorly drained and with a natural high water table.

## General land use suitability ratings

### Awaiti sand

Land use	Suitability rating	Management considerations
Arable	Moderate to low	Low fertility, summer droughts, fragile soil structure.
Horticulture	Moderate to low	Low fertility, summer droughts, fragile soil structure, fluctuating ground water table.
Intensive pasture	Moderate	Summer droughts, low fertility.
Forestry	Moderate	Fluctuating ground water table.

### Management practices to improve suitability

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Awaiti sand** (Note subsoil mottling and thin layer of Kaharoa Tephra at 40 cm depth)

## Soil Series Name: Awakaponga (Ag)

### Overview

Awakaponga soil series occur on the flat drained back swamp lowlands associated with the Rangitaiki River. They are derived from fine alluvium derived from rhyolitic tephra and minor weathered greywacke, frequently interlayered with peaty material. The soils are typically layered with several buried topsoils resulting from previous flooding. Strong brown mottling and staining occurs below the topsoil. Peaty layers are common. The soils are classified as **Mottled-acidic Fluvial Recent Soils**. Present land use includes dairying, maize cropping and some urban use.

### Physical properties

**Texture:** Silt loam over peat and sand

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** 60 – 100 cm

**Rooting barrier:** Fluctuating water table

**Drainage class:** Imperfectly to poorly drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** Moderate to high (142 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (85 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.38 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 8.6%

**Topsoil P retention:** Medium (33%)

**Available P, Ca, Mg and K:** Low to medium

### Soil types/variations

**Awakaponga silt loam** and **Awakaponga silt loam on peat** have been recognised.

#### ***Associated and similar soils***

Inclusions of **Paroa series** may occur. **Rangitaiki series** are well drained and occur closer to the riverbed.

## ***General land use suitability ratings***

### **Awakaponga silt loam on peat and Awakaponga silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate to low	Imperfect to poor drainage, structural vulnerability.
Horticulture	Moderate to low	Imperfect to poor drainage, structural vulnerability.
Intensive pasture	Moderate	Imperfect to poor natural drainage. high water table in wet winter.
Forestry	Low	Imperfect to poor natural drainage.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for cropping or market gardening.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Employ surface drainage.



**Awakaponga silt loam**

## Soil Series Name: Awakeri (Aw)

### Overview

Awakeri soil series occur on the Rangitaiki Plains around Western Drain Road on flat basins and former stream ridges. They are derived from very thin Tarawera Tephra and Kaharoa Tephra, on very thin peat and alluvium derived from Taupo Tephra. Profiles have black friable sand on very dark brown gravelly sand and pale yellow loose sand. These rest on some 10 cm dark reddish-brown peat overlying brown coarse sand. The soils are classified as **Typic Perch-gley Pumice Soils**. Present land use consists of dairying, dry stock and some horticulture (strawberries).

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 5 – 10%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Imperfectly drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** High (151 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (78 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 13.8%

**Topsoil P retention:** Moderate (40%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Awakeri sand** is the dominant soil type. Some variation in thickness of tephra layers, alluvium and peat occurs.

#### ***Associated and similar soils***

**Te Teko series** have coarser textured subsoils while **Pongakawa series** occur on adjacent peaty basins.

## **General land use suitability ratings**

### **Awakeri sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Medium	Weakly developed topsoil structure, low fertility, imperfect drainage.
Horticulture	Medium	Weakly developed topsoil structure, low fertility, imperfect drainage.
Intensive pasture	High	Low fertility, imperfect drainage.
Forestry	Not suitable	Imperfect drainage.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for cropping or market gardening.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Employ artificial drainage (open drains).



**Awakeri sand** (Note Kaharoa Tephra at 40 to 50 cm depth)

## Soil Series Name: Galatea (G)

### Overview

Galatea soil series occur in the Galatea Basin on flat to undulating surfaces. Parent materials consist of 60 cm Kaharoa Tephra, on 80 cm Taupo Tephra on Whakatane Tephra. Typically the soils have friable black sandy topsoils overlying brown coarse sand and pale yellow coarse sand. The Taupo buried topsoil is dark yellowish brown sandy loam which rests on pale yellow fine pumice gravel (Taupo lapilli). The soils are classified as **Immature Orthic Pumice Soils**. The soils are used for dairying, dry stock and some fodder cropping.

### Physical properties

**Texture:** Sand over loam

**Topsoil clay content:** 3 – 8%

**Potential rooting depth:** Unlimited

**Rooting barrier:** None

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (102 mm)

**Profile readily available water (0 – 100 cm):** Moderate (64 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical Properties

**Topsoil organic matter:** 8.6 – 12%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Galatea sand** is the main soil type with very little variation. Small areas have subsoil mottling in shallow depressions.

#### ***Associated and similar soils***

The soils grade into **Kopuriki series** towards the north of the Galatea Basin where thin Tarawera Tephra overlies the Galatea soil profile. **Horomanga series** occur on gently sloping fans in eastern parts of Galatea Basin. **Rangitaiki series** are alluvial soils along the Rangitaiki River that flows through the Basin.


## **General land use suitability ratings**

### **Galatea sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Droughty soil, weak topsoil structure; low fertility, wind erosion if cultivated during dry spells.
Horticulture	Low	Cool climate, droughty soil, weak topsoil structure low fertility, wind erosion if cultivated during dry spells.
Intensive pasture	Low to moderate	Summer droughts, low fertility.
Forestry	Moderate to high	Droughts affecting young trees.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.
- Irrigation is used extensively on Galatea soils.
- Forestry: avoid windrowing and damage to topsoils.



**Galatea sand** (scale in m)

## Soil Series Name: Hanaia (Hi)

### Overview

Hanaia soil series occur on flat to undulating broad valley floors, chiefly in the Opotiki area. The soils are derived from alluvium derived from rhyolitic tephra that eroded off valley sides. A thin layer of Taupo Pumice may occur in some areas. Profiles consist of very dark greyish brown sandy loam overlying mottled sandy loam and silt loam. Soil classification is **Typic Orthic Gley Soil**. Land use is dairying, dry stock and cropping (maize).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 15 – 20%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier

**Drainage class:** Imperfectly to poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** High (157 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (95 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 9.6 – 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Hanaia silt loam** and **Hanaia sandy loam** have been mapped. Thin peaty layers occur in Opotiki near the boundary with Paerata series.

### *Associated and similar soils*

**Otara, Apanui and Waioeka series** on the Opotiki Plains are derived from alluvium from greywacke and rhyolitic tephra and are mostly heavier textured with thinner tephra layers. **Taho series** are on fans derived from tephra only. **Paerata series** are on valley floors with peaty layers.

### **General land use suitability ratings**

#### **Hanaia silt loam and Hanaia sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate to low	Low fertility, imperfect to poor natural drainage, fragile soil structure.
Horticulture	Low	Low fertility, imperfect to poor natural drainage, fragile soil structure.
Intensive pasture	Moderate	Imperfect to poor natural drainage.
Forestry	Unsuitable	Imperfect to poor natural drainage.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Employ surface drainage.
- Plant green crops to increase organic matter and thicken topsoils.



**Hanaia silt loam** (Note subsoil mottling)

## Soil Series Name: Horomanga (Hm)

### Overview

Horomanga soil series occur on flat to slightly sloping fans on the eastern side of the Galatea Basin. The soils are derived from alluvium and colluvium from tephra and greywacke. Soil profiles are variable but generally consist of very dark brown sand overlying pale yellow coarse sand. The soils are classified as **Typic Tephric Recent Soils**. Land use is mostly dairying or dry stock.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 2 – 6%

**Potential rooting depth:** Unlimited

**Rooting barrier:** None within 1 m depth

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (97 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (75 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.38 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** Less than 8.6 to 12%

**Topsoil P retention:** Medium (32%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Horomanga sand** is the main soil type. **Horomanga mottled silt loam** (poorly drained) occurs at the base of the fans.

#### *Associated and similar soils*

**Galatea series** are on flat terrace-like surfaces derived from airfall Kaharoa and Taupo Tephra. **Kopuriki series** are like Galatea series but with a thin (more than 7 cm) layer of Tarawera Tephra on top. **Rangitaiki series** are alluvial soils along the Rangitaiki River.

## **General land use suitability ratings**

### **Horomanga sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Fragile soil structure, low fertility, summer droughts.
Horticulture	Low	Cool climate, fragile soil structure, low fertility, summer droughts.
Intensive pasture	Low to moderate	Summer droughts, low fertility.
Forestry	Low to moderate	Summer droughts, low fertility.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.
- Forestry: avoid windrowing and damage to topsoils.



**Horomanga sand**

## Soil Series Name: Iwiroa (Ir)

### Overview

Iwiroa soil series occur on slightly sloping fans in the Eastern Bay of Plenty. The soils are derived from colluvium from tephra and greywacke. Soil profiles are variable but generally consist of very dark grey silt loam overlying greyish-brown sandy loam and light brownish-grey sandy clay loam. Angular greywacke fragments are common. The soils are classified as **Typic Orthic Gley Soils**. Land use is mostly dairying or dry stock.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** 60 – 80 cm

**Rooting barrier:** Wet subsoil

**Drainage class:** Poorly drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** Moderate (118 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (75 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** Less than 6.9 to 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Iwiroa silt loam** is the main soil type. Buried topsoils occur locally as a result of recent erosion and accumulation. The upper part of the fan is generally drier and the foot of the fan wetter.

#### ***Associated and similar soils***

**Tutaetoko series** are on adjacent steep slopes while **Opotiki series** are on adjacent hilly slopes.

## General land use suitability ratings

### Iwiroa silt loam

Land use	Suitability rating	Management considerations
Arable	Low	Poor drainage, fragile soil structure, sometimes stony soils, low fertility.
Horticulture	Not recommended	Poor drainage, fragile soil structure, sometimes stony soils, low fertility.
Intensive pasture	Low to moderate	Poor drainage, low fertility.
Forestry	Not recommended	Poor drainage.

### Management practices to improve suitability

- Employ surface drainage.
- Use zero or minimum tillage methods for fodder cropping.
- Plant green crops to increase organic matter and thicken topsoils.



**Fan with Iwiroa soils extending on river flats. Waipua stony sand in the foreground around the streambed. Owaka soils on the next terrace. Iwiroa soils on the fan with the rushes.**

## Soil Series Name: Kawerau (Kr)

### Overview

Kawerau soil series occur on the southern part of the Rangitaiki Plains on flat to undulating terraces. They are derived from thin Tarawera Tephra on pumice alluvium derived from Kaharoa and Taupo Tephra. Profiles show very dark greyish-brown coarse sand on dark brown sandy gravel. This rests on dark grey and brown loose sand. The soils are classified as **Immature Orthic Pumice Soils**. These are the first soils to dry out on the Rangitaiki Plains. Present land use includes dry stock, some dairying and some pip fruit orchards.

### Physical properties

**Texture:** Skeletal

**Topsoil clay content:** 1 – 5%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant root barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (37 mm)

**Profile readily available water (0 – 100 cm):** Low (28 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Kawerau sand** is generally well drained while **Kawerau mottled gravelly sand** has mottling in the subsoil in local shallow depressions and is regarded as imperfectly drained.

### *Associated and similar soils*

**Tarawera series** have thicker and coarser Tarawera Tephra (more than 20 cm). **Te Teko series** have finer textures.

## **General land use suitability ratings**

### **Kawerau sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Low fertility, summer droughts, fragile soil structure.
Horticulture	Low	Low fertility, summer droughts, fragile soil structure.
Intensive pasture	Moderate to low	Summer droughts, low fertility, good winter grazing.
Forestry	Moderate	Summer droughts, flattish areas better used for pasture.

### **Kawerau mottled gravelly sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Low fertility, summer droughts, fragile soil structure.
Horticulture	Low to moderate	Fluctuating ground water levels, summer droughts.
Intensive pasture	Moderate to low	Summer droughts, low fertility, good winter grazing.
Forestry	Moderate to low	Fluctuating ground water levels, summer droughts, flattish areas better used for pasture.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods with cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Kawerau sand**

## Soil Series Name: Kopeopeo (Koe)

### Overview

Kopeopeo soil series have been mapped in the northern part of Whakatane District, on the Rangitaiki Plains on inland dunes and on subdued ridges parallel to the present shoreline. They are derived from thin to very thin Tarawera, Kaharoa, Taupo, Waimihia and Whakatane Tephra on wind-blown sand. Soil profiles show black coarse loamy sand on pale yellowish-brown and brownish-yellow sand. An iron/humus pan occurs at about 80 to 110 cm depth. The soils are classified as **Humus-pan Pan Podzols**. Current land use includes dairying, dry stock and some urban use in Whakatane.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 4 – 8%

**Potential rooting depth:** 80 to 100 cm

**Rooting barrier:** Humus/iron pan at 80 to 100 cm depth

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (80 mm)

**Profile readily available water (0 – 100 cm):** Moderate (60 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 20.7%

**Topsoil P retention:** High (60%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Kopeopeo loamy sand** is the predominant soil type. An unmapped peaty phase occurs in swales where wind-blown sand is interlayered with peat. Inclusions consist of areas with less tephra after wind erosion or human alterations.

#### ***Associated and similar soils***

**Kairua series** in the Te Puke area with less tephra and the iron/humus pan generally closer to the surface; **Piripai series** without Taupo Tephra occurring closer to the shore; **Pikowai series** on the fore dunes without tephra; and **Te Rahu series** on older inland dunes where thicker tephra overlies wind-blown sand.

## **General land use suitability ratings**

### **Kopeopeo loamy sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low to not suitable	Coarse textures, summer droughts. low fertility, fragile topsoil structure susceptible to wind erosion if cultivated during dry spells.
Horticulture	Low to not suitable	Coarse textures, summer droughts, low fertility, fragile topsoil structure susceptible to wind erosion if cultivated during dry spells.
Intensive pasture	Low	Low fertility, summer droughts.
Forestry	Moderate	Low fertility, low soil moisture at establishment.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods when working the soil.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Use for winter grazing during wet periods.



**Kopeopeo loamy sand**

## Soil Series Name: Kopuriki (Kpu)

### Overview

Kopuriki soil series have been mapped in the northern Galatea Basin on flat to undulating surfaces. Soils are formed from more than 7 cm Tarawera Tephra, on Kaharoa, Taupo and Whakatane Tephra layers. Soil profiles show very dark greyish brown sandy gravel (Tarawera) on black firm sand (Kaharoa). This overlies pumice gravel layers (Kaharoa and Taupo lapilli) and yellowish-brown sandy loam (Whakatane Tephra). The soils are classified as **Immature Orthic Pumice Soils**. Current land use includes dry stock and dairying.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 4 – 6%

**Potential rooting depth:** Unlimited

**Rooting barrier:** None within 1 m depth

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (82 mm)

**Profile readily available water (0 – 100 cm):** Moderate (54 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Kopuriki gravelly sand** is the main soil type.

#### ***Associated and similar soils***

**Galatea series** where Tarawera Tephra is less than 7 cm thick; **Matahina series** on rolling and hilly surfaces with less than 20 cm coarse Tarawera lapilli at the surface

## **General land use suitability ratings**

### **Kopuriki gravelly sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Low fertility, summer droughts, fragile topsoil structure susceptible to wind erosion if cultivated during dry spells.
Horticulture	Low to not suitable	Cool climate, low fertility, summer droughts, fragile topsoil structure susceptible to wind erosion if cultivated during dry spells.
Intensive pasture	Moderate to low	Low fertility, summer droughts.
Forestry	Moderate	Flattish land better used for pastoral uses, low fertility, fragile topsoil.

### **Management practices to improve suitability**

- Use zero or minimum tillage methods when working the soil.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Restrict topsoil damage during harvesting.



**Kopuriki gravelly sand**

## Soil Series Name: Kukumoa (Ku)

### Overview

Kukumoa soil series occur in moderate areas west of Opotiki and smaller areas west of the Waioeka River and Waiotahi River on imperfectly drained parts of low-lying river flats. Soil profiles show shallow brown silt loam topsoils overlying olive mottled fine sandy loam. Buried topsoils from past flooding are common. The soils are classified as **Mottled Fluvial Recent Soils**. Current land use is dairying, dry stock and cropping (maize).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 15 – 20%

**Potential rooting depth:** 30 – 50 cm

**Rooting barrier:** Fluctuating ground water table

**Drainage class:** Imperfectly drained with fluctuating ground water table

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** Moderate to high (95 mm)

**Profile readily available water (0 – 100 cm):** Moderate (57 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 8.6%

**Topsoil P retention:** Medium (33%)

**Available Ca, Mg and K:** High P; High to very high Ca, Mg, K

### Soil Types/Variations

Both **silt loam** and **sandy loam** topsoils occur.

#### ***Associated and similar soils***

**Amokura series** are poorly drained and occur in former oxbows or shallow depressions. **Rangitaiki series** and **Ruatoki series** are well drained soils of the same river flat and occur adjacent to the river. **Opouriao series** are well drained soils of the levee above the aforementioned soils.

## **General land use suitability ratings**

### **Kukumoa silt loam and sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Imperfect drainage, frequent flooding if not adequately protected by stop banks.
Horticulture	Moderate	Imperfect drainage, frequent flooding if not adequately protected by stop banks.
Intensive pasture	High	Imperfect drainage, frequent flooding if not adequately protected by stop banks.
Forestry	Not suitable	Imperfect drainage.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods when working the soil.
- Employ surface drainage.



**Kukumoa silt loam** (Note mottling in the subsoil)

## Soil Series Name: Manawahe (Mj)

### Overview

Manawahe soil series occur west of the Rangitaiki Plains and east of the Rangitaiki River on rolling to hilly terrain. The soils are formed from 7 to 20 cm sandy Tarawera Tephra on Kaharoa and Taupo Tephra overlying Whakatane Tephra and older weathered rhyolitic tephra. Soil profiles show dark brown gritty sand on brown and light olive brown sand overlying pale yellow fine pumice gravel and yellowish-brown loamy sand. The soils are classified as **Immature Orthic Pumice Soils**. Current land use includes dairying, dry stock, some fodder cropping, and forestry (*Pinus radiata*).

### Physical properties

**Texture:** Sand and sand over skeletal

**Topsoil clay content:** 2 – 3%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (83 mm)

**Profile readily available water (0 – 100 cm):** Moderate (62 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Manawahe loamy sand** occurs on flat to rolling land and **Manawahe hill soils** are on hilly slopes. Thickness of Tarawera Tephra varies from 7 to 20 cm. Locally, slightly reddish subsoil occurs.

#### ***Associated and similar soils***

**Matahina series** occur on rolling to hilly land with less than 20 cm gravelly Tarawera Tephra on Kaharoa Tephra. They have less water-holding capacity than Manawahe series because of coarser texture. **Pukemaku series** are on steep and very steep slopes where shallower tephra overlies ignimbrite. **Tarawera series** are on easy rolling to hilly land. More than 20 cm gravelly Tarawera Tephra overlies Kaharoa Tephra. They have less water-holding capacity than Manawahe series because of coarser texture. **Whakatane series** are on easy rolling to hilly land with very thin sandy Tarawera Tephra and Kaharoa Tephra overlying weathered rhyolitic tephra.


## **General land use suitability ratings**

### **Manawahe loamy sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low to moderate (hill soils not suitable)	Low fertility, fragile topsoil.
Horticulture	Low to moderate (Hill soils not suitable)	Low fertility, fragile topsoil.
Intensive pasture	Moderate	Low fertility, fragile topsoil especially on hilly slopes.
Forestry	High	Low fertility.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods when working the soil.
- Avoid cultivating the soil during dry spells.



**Manawahe loamy sand** (Note deep topsoil formed from fine Tarawera Tephra and the buried topsoil of Kaharoa Tephra, scale in ft)

## Soil Series Name: Mangaomeko (MaH)

### Overview

Mangaomeko soil series occur in hill country south of Cape Runaway on moderately steep slopes under 1200 to 1800 mm annual rainfall. The soils are formed from mudstone and profiles have black silt loam topsoils overlying olive yellow loam or silt loam with an increasing amount of weathered mudstone fragments with increasing depth. Brownish-yellow weathered mudstone generally occurs at 70 cm depth. The soils are classified as **Typic Orthic Brown Soils**. Current land use is dry stock.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 30%

**Potential rooting depth:** 50 to 90 cm

**Rooting barrier:** Weathering rock

**Drainage class:** Well drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** Moderate (95 mm)

**Profile readily available water (0 – 100 cm):** Moderate (53 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.53 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 5.2 – 11.2%

**Topsoil P retention:** Medium (36%)

**Available P, Ca, Mg and K:** Probably medium levels

### Soil types/variations

**Mangaomeko hill soils** are quite variable. Deeper and shallower soil profiles occur frequently. Small pockets of tephra are on stable parts of the hill country.

#### ***Associated and similar soils***

**Tuparoa series** occur on adjacent steep and very steep slopes with generally shallower soil profiles on mudstone.

## **General land use suitability ratings**

### **Mangaomeko hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, severe erosion potential.
Horticulture	Not suitable	Hilly slopes, severe erosion potential.
Intensive pasture	Low	Hilly slopes, severe erosion potential.
Forestry	Low to moderate	Hilly slopes, moderate erosion potential.

### ***Management practices to improve suitability***

- Intensive land use would accelerate creep and slip erosion; therefore, low stocking rates would be advisable.



**Mangaomeko hill soil**

## Soil Series Name: Man-made (MM)

### Overview

Man-made soils are soils that have been altered by humans. These soils are extremely variable and no detailed morphological description has been given here. Generally, parent materials are tephra. Large areas are shown on soil maps, but many smaller areas are not. The soils are classified as **Mixed Anthropic Soils** in the New Zealand Soil Classification. Recontoured land supports kiwifruit in coastal areas.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 - 30%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** High (170 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (88 mm)

**Topsoil bulk density:** 1.41 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** Not tested

**Topsoil P retention:** Not known

**Available P, Ca, Mg and K:** Probably low levels

### Soil types/variations

Man-made soils vary from large earthworks to recontoured land.

### *General land use suitability ratings*

#### Man-made soils

Land use	Suitability rating	Management considerations
Arable	Low	Not considered.
Horticulture	Low to moderate	Recontoured land planted in kiwifruit.
Intensive pasture	Low to moderate	Depending on adequate restoration.
Forestry	Moderate	Depending on adequate restoration.

### ***Management practices to improve suitability***

- Replacement of adequate topsoil.
- Ripping and restoring subsoils and thus removing root barriers.



**Man-made soil**

## Soil Series Name: Mataheiiia (Map)

### Overview

Mataheiiia soil series occur on flat river levees along the main rivers of Cape Runaway. They are formed from alluvium derived from greywacke, sandstone, mudstone and tephra. Profiles show very dark greyish-brown silt loam topsoils overlying olive brown silt loam and light yellowish brown sand. The soils are classified as **Weathered Fluvial Recent Soils**. Land use on improved pasture is dairying, dry stock and also maize.

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 20 – 30%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to high (139 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (99 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 to 8.6%

**Topsoil P retention:** Low (19%)

**Available P, Ca, Mg and K:** Low to moderate

### Soil types/variations

**Matehiiia silt loam** is the main soil type. Inclusions with faint mottling below 60 cm depth also occur.

#### ***Associated and similar soils***

**Oweka series** are on slightly lower river terraces with paler (less organic matter) topsoils and subject to more frequent flooding where not protected by stop banks. **Opouriao series** in the Central Bay of Plenty are derived from slightly different parent materials.

## **General land use suitability ratings**

### **Mateheia silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate to high	Fragile structure, slight flooding risk.
Horticulture	Moderate to high	Fragile structure, slight flooding risk.
Intensive pasture	Moderate to high	Fragile structure, slight flooding risk.
Forestry	Low	Slight flooding risk, fragile structure.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods when working the soil.
- Sow green crops to thicken and improve topsoil.



**Mataheia silt loam** (Note: Spade 90 cm long)

## Soil Series Name: Matahina (Mb)

### Overview

Matahina soil series occur in central Whakatane District in the Matahina Forest locality. The soils are formed from less than 20 cm gravelly Tarawera Tephra, on Kaharoa Tephra, on Taupo Tephra and weathered rhyolitic tephra. Topsoils are black gravelly and sand overlying pale yellow coarse sand (Kaharoa lapilli) which rest on very dark brown greasy sandy loam and pale yellow pumice lapilli layer (Taupo lapilli). The soils are classified as **Immature Orthic Pumice Soils**. Current land use is forestry (*Pinus radiata*) and orchards near the Rangitaiki Plains where the climate is milder.

### Physical properties

**Texture:** Sand (skeletal on hilly slopes)

**Topsoil clay content:** 0 – 4%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (74 mm)

**Profile readily available water (0 – 100 cm):** Low (43 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Matahina gravelly sand** occurs on flat to rolling land and **Matahina hill soils** on hilly slopes. Thickness of Tarawera Tephra varies from 7 to 20 cm, Kaharoa Tephra from 50 to 100 cm, and Taupo Tephra from 40 to 60 cm. Many shallow slips and creep erosion occur on hilly slopes (movement of Tarawera lapilli downslope especially in pasture).

#### ***Associated and similar soils***

**Manawahe series** occur on rolling to hilly land with less than 20 cm sandy Tarawera Tephra on Kaharoa Tephra. These soils have somewhat higher water-holding capacity than Matahina series because of finer textures.

**Pukemaku series** occur on steep and very steep slopes where shallower tephra overlies ignimbrite.

**Tarawera series** are on easy rolling to hilly land where more than 20 cm gravelly Tarawera Tephra overlies Kaharoa Tephra. The soils have lower water-holding capacity than Manawahe series because of coarser textures.

**Whakatane series** occur on easy rolling to hilly land where very thin sandy Tarawera Tephra and Kaharoa Tephra overly weathered rhyolitic tephra.

### **General land use suitability ratings**

#### **Matahina gravelly sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low to not suitable	Low fertility, fragile topsoil, low water-holding capacity, creep erosion.
Horticulture	Not suitable	Cool climate, creep erosion, rolling topography, infertile soils.
Intensive pasture	Moderate to low	Low fertility, fragile topsoil, creep erosion on rolling slopes, low water-holding capacity.
Forestry	Moderate	Low fertility, creep erosion on rolling slopes.

#### **Matahina hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, severe erosion potential, low water-holding capacity, low fertility, fragile topsoil, creep erosion.
Horticulture	Not suitable	Hilly slopes, cool climate.
Intensive pasture	Low	Hilly slopes, severe erosion potential, low water-holding capacity, low fertility, fragile topsoil, creep erosion.
Forestry	Moderate	Low fertility, severe erosion potential.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods when working the soil.
- Avoid cultivating the soil during dry spells.
- Avoid extensive damage to topsoils during harvesting.



**Matahina gravelly sand**

## Soil Series Name: Matakaoa (Mak)

### Overview

Matakaoa soil series occur on flat to undulating terrace country and on hilly slopes at Cape Runaway area. The soils are formed from rhyolitic tephra on loess and basalt. Soil profiles consist of very dark brown sandy loam, grading to dark brown sandy loam, which rest on yellowish-brown and brownish-yellow loam and silt loam. The soils are classified as **Typic Orthic Allophanic Soils**. Land use consists of dry stock, dairying and some maize cropping.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 10 – 18%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** High (167 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (95 mm)

**Topsoil bulk density:** 0.78 g/cm<sup>3</sup>

**Subsoil bulk density:** 0.86 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 12.9 – 22.4%

**Topsoil P retention:** High (83%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Matakaoa sandy loam** is located on flat to rolling terraces and **Matakaoa hill soils** are on hilly slopes. The latter has shallower tephra on loess and basalt.

#### ***Associated and similar soils***

**Wharekahika series** have mottling in the subsoil; **Potikirua series** are on steep and very steep slopes, derived from basalt with very little tephra; and **Te Kaha series** which also formed from rhyolitic tephra but contains lesser allophane (**Typic Allophanic Brown Soils**).

## **General land use suitability ratings**

### **Matakaoa sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Low fertility, fragile topsoil, some rolling slopes.
Horticulture	Moderate	Low fertility, fragile topsoil, some rolling slopes.
Intensive pasture	Moderate to high	Low fertility, fragile topsoil.
Forestry	Moderate to high	Low fertility.

### **Matakaoa hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, erosion potential, creep erosion.
Horticulture	Not suitable	Hilly slopes, erosion potential, creep erosion.
Intensive pasture	Moderate	Hilly slopes, erosion potential, creep erosion.
Forestry	Moderate to high	Low fertility, erosion potential, hilly slopes.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods when working the soil.
- Avoid cultivating the soil during dry spells.
- Avoid extensive damage to topsoils during harvesting.



**Matakoa sandy loam**

## Soil Series Name: Matao (Ma)

### Overview

Matao soil series occur on valley floors in the Opotiki region. The soils are formed from very thin alluvium, on very thin Kaharoa and Taupo Tephra (two distinct sandy layers), on peat. In places, woody materials occur in the subsoil and such soils have been mapped separately. A common soil profile is black peaty silt loam on light grey fine sand (Kaharoa Tephra) on very dark brown peat with a light yellowish brown sand layer (Taupo Tephra) at 40 cm depth. The soils are classified as **Mellow Mesic Organic Soils**. Land use is pasture for dairying and/or dry stock. Blueberries could be grown if ground water levels can be strictly controlled.

### Physical properties

**Texture:** Loamy peat over sand

**Topsoil clay content:** 8 – 9%

**Potential rooting depth:** 50 – 70 cm

**Rooting barrier:** Fluctuating ground water levels (50 to 70 cm approximately)

**Drainage class:** Poorly drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** Moderate to low (86 mm)

**Profile readily available water (0 – 100 cm):** Moderate (70 mm)

**Topsoil bulk density:** 0.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 0.18 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 34.5 - 69%

**Topsoil P retention:** High (70%) in peaty horizons but low to very low in all other horizons

**Available P, Ca, Mg and K:** Low levels

### Soil types/variations

**Matao peaty silt loam.** Inclusions of **Matao peaty silt loam, woody subsoil phase** with old stumps in the lower subsoil. Wood remnants may occur close to the surface with over drainage.

#### *Associated and similar soils*

**Paerata series** with thin peat layers; **Hanaia series** without peat; and **Taho series** on former slumped areas usually near the head of the valleys.

## ***General land use suitability ratings***

### **Matao peaty silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Poor drainage, fragile topsoils.
Horticulture	Moderate	Poor drainage.
Intensive pasture	Moderate	Poor drainage, pugging under heavy stocking.
Forestry	Not suitable	Poor natural drainage.

### ***Management practices to improve suitability***

- Employ surface drainage taking care not to over-drain due to the risk of peat shrinkage.
- Remove stumps from the surface.



**Matao peaty silt loam** (Note Kaharoa Tephra at 20 cm and Taupō Tephra at 40 to 60 cm depth)

## Soil Series Name: Matatā (Maa)

### Overview

Matata soil series occur in the western part of the Rangitaiki Plains on active low-angled fans. Parent materials are fine pumiceous colluvium with minor greywacke and locally strongly weathered tephra. At Matata locality, the subsurface is wind-blown sand. Soil profiles have thick (26 cm) very dark grey sandy loam, resting on very dark grey and brown gravelly sand. The soils are classified as **Mottled Tephric Recent Soils**. Current land use is dry stock.

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 10 – 20%

**Potential rooting depth:** 50 – 80 cm

**Rooting barrier:** Fluctuating ground water levels in some parts

**Drainage class:** Imperfectly drained to poorly drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (117 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (97 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.38 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (32%)

**Available P, Ca, Mg and K:** Medium to high

### Soil types/variations

**Matata sandy loam** is the main soil type for this series.

#### ***Associated and similar soils***

**Opouriao series, Paroa series, and Te Rahu series** all occur at the toes of the fans. **Whakatane series, Manawahe series and Matahina series** occur above the fans.

## **General land use suitability ratings**

### **Matata sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Imperfect to poor natural drainage.
Horticulture	Low	Imperfect to poor natural drainage.
Intensive pasture	Moderate	Imperfect to poor drainage, pugging under heavy stocking.
Forestry	Not suitable	Imperfect to poor natural drainage.

### ***Management practices to improve suitability***

- Employ surface drainage.
- Stabilise areas above the fan so no further soil accumulation can occur.



**Matata sandy loam**

## Soil Series Name: Matawai (Mx)

### Overview

Matawai soil series occur in eastern parts of Whakatane District in Urewera National Park on rolling and hilly slopes under 2000 to 2500 mm annual rainfall at elevations above 700 m a.s.l. Parent materials are very thin Kaharoa and Taupo Tephra on Waimihia Tephra and weathered rhyolitic tephra overlying greywacke. Soil profiles have thin reddish-brown loamy sand on greyish-brown sand (hard, massive) resting on reddish brown and yellowish-red pumice gravel and dark yellowish-brown loamy sand and sandy loam. The soils are classified as **Placic Pan Podzols**. Land use is protection forest (mainly beech) with some cut-over bush.

### Physical properties

**Texture:** Loam over sand (if sandy loam); Sand (if hill soil)

**Topsoil clay content:** 10 – 15% (if sandy loam); 3 – 4% (if hill soil)

**Potential rooting depth:** 20 – 30 cm (if sandy loam); 60 – 70 cm (if hill soil)

**Rooting barrier:** Iron pan

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (39 – 45 mm)

**Profile readily available water (0 – 100 cm):** Low (32 – 36 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 20.7%

**Topsoil P retention:** Medium (44%)

**Available P, Ca, Mg and K:** Low levels

### Soil types/variations

**Matawai sandy loam** are on flat to rolling land and **Matawai hill** soils on hilly slopes. Kaharoa and Taupo Tephra are usually thicker on foot slopes.

#### ***Associated and similar soils***

**Urewera series** are on steep and very steep slopes with thinner tephra layer overlying greywacke. **Ruakituri series** occur at lower elevation and somewhat lower annual rainfall. They do not have an iron pan.

## **General land use suitability ratings**

### **Matawai sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Unsuitable	Cold climate, very strong leaching, low fertility, fragile topsoil.
Horticulture	Unsuitable	Cold climate, very strong leaching, low fertility, fragile topsoil.
Intensive pasture	Low to unsuitable	Cold climate (short growing season), very strong leaching, very low fertility.
Forestry	Low	Cold climate, low fertility, strong leaching.

### **Matawai hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Cold climate, hilly slopes, very strong leaching, low fertility, fragile topsoil.
Horticulture	Not suitable	Cold climate, hilly slopes, very strong leaching, low fertility, fragile topsoil.
Intensive pasture	Low to not suitable	Cold climate (short growing season), very strong leaching, very low fertility, hilly slopes.
Forestry	Low	Cold climate, low fertility, strong leaching, hilly slopes, fragile topsoil.

### ***Management practices to improve suitability***

- Protection forest is the best option.



**Matawai sandy loam**

## Soil Series Name: Matuku (Mtk)

### Overview

Matuku soil series occur in the western part of the Rangitaiki Plains with small areas in Onepu, Braemar Road, Hawkins Junction and Awaiti locations, on flat former flood plains of the Tarawera River. Parent materials of the soil include diatomaceous earth, peat and pumice alluvium with a very thin cover of Tarawera Tephra. Soil profiles have very dark greyish-brown fine sandy loam with coarse sand (Tarawera lapilli) and fine yellowish-brown mottles. They rest on light brownish-grey silt, on very dark brown peaty silt loam overlying pale brown sandy loam and sand. The soils are classified as **Peaty Orthic Gley Soils**. Current land use includes dry stock, dairying and breeding stock.

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 15 – 20%

**Potential rooting depth:** 50 – 80 cm

**Rooting barrier:** Fluctuating ground water levels

**Drainage class:** Poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Moderate (98 mm)

**Profile readily available water (0 – 100 cm):** Moderate (65 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** Medium

### Soil types/variations

**Matuku fine sandy loam** is the main soil type.

#### *Associated and similar soils*

**Onepu series** have coarser textures and without diatomaceous layers. **Rangitaiki series** are on gravelly ridges.

## **General land use suitability ratings**

### **Matuku fine sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Poor natural drainage, fluctuating ground water levels, fragile soil structure.
Horticulture	Low	Poor natural drainage, fluctuating ground water levels, fragile soil structure.
Intensive pasture	Moderate	Poor natural drainage, pugging under heavy stocking.
Forestry	Not suitable	Poor natural drainage.

### ***Management practices to improve suitability***

- Employ surface drainage.
- Plant green crops to build up organic matter in the topsoil.



**Matuku silt loam**

## Soil Series Name: Mōtū (MotH)

### Overview

Motu soil series occur on hilly slopes in coastal areas and in the Raukumara Ranges of the Eastern Bay of Plenty under 1400 to 2000 mm annual rainfall. Parent material is greywacke. Profiles show brown silt loam overlying dark yellowish-brown gritty loam. Angular greywacke fragments increase with increasing depth. The soils are classified as **Typic Orthic Brown Soils**. Much of the land cover is indigenous forest with some areas in pasture (dry stock).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 22 – 27%

**Potential rooting depth:** 25 – 50 cm

**Rooting barrier:** Fractured rock

**Drainage class:** Well drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** Low (35 mm)

**Profile readily available water (0 – 100 cm):** Very low (24 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.53 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 5.2 – 11.2%

**Topsoil P retention:** Medium (36%)

**Available P, Ca, Mg and K:** Not known

### Soil types/variations

**Motu hill soils** can have common shallow soil profiles occurring where only 20 cm or so weathered material overlies shattered greywacke.

#### ***Associated and similar soils***

**Raukumara series** on steep and very steep slopes with often shallow soil profiles.

## **General land use suitability ratings**

### **Motu hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, shallow soil profiles, low water-holding capacity.
Horticulture	Not suitable	Hilly slopes, shallow soil profiles, low water-holding capacity.
Intensive pasture	Low to not suitable	Hilly slopes, shallow soil profiles, low water-holding capacity.
Forestry	Not suitable	Hilly slopes, shallow soil profiles, low water-holding capacity.

### ***Management practices to improve suitability***

- Prevent erosion by keeping vegetation on the slopes as much as possible.



**Motu hill soil**

## Soil Series Name: Ngatiawa (NS)

### Overview

Ngatiawa soil series occur on steep and very steep slopes in north-eastern parts of Whakatane District. Parent materials are very thin Kaharoa and Taupo Tephra on Whakatane Tephra and older rhyolitic tephra overlying sandstone. Soils profiles show very dark greyish brown fine sandy loam, on olive brown sandy loam and light yellowish-brown firm fine sandy loam, resting on light yellowish brown firm weathered sandstone. The soils are classified as **Typic Tephric Recent Soils**. Land use on pasture is dry stock (sheep) but large areas are in shrub (manuka).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 15 – 20%

**Potential rooting depth:** 40 – 80 cm

**Rooting barrier:** Weathered rock

**Drainage class:** Well drained

**Permeability:** Rapid over moderate

**Profile total available water (0 – 100 cm):** Moderate to low (77 mm)

**Profile readily available water (0 – 100 cm):** Low (49 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (32%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Ngatiawa stepland soils** have thinner tephra which occurs on eroded slopes.

#### *Associated and similar soils*

**Whakatane series** occur on easy rolling to hilly slopes with thicker tephra layers. **Tawhia series** are found on steep and very steep slopes where tephra overlies greywacke.

## **General land use suitability ratings**

### **Ngatiawa steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep and very steep slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low	Steep and very steep slopes, low fertility, risk of accelerated erosion.
Forestry	Low	Steep and very steep slopes, accelerated erosion risk, low fertility.

### **Management practices to improve suitability**

- Avoid high stocking rates.
- Forestry: avoid damage to topsoils.



**Ngatiawa steepland soil landscape**

## Soil Series Name: Ōhope (Oe)

### Overview

Ohope soil series occur on coastal fore dunes in Western Bay of Plenty and East of Whakatane. The soils are formed from wind-blown sand. Soil profiles are weakly developed consisting of 10 cm light brownish-grey sand on light grey sand. They are classified as **Sandy Raw Soils**. Land use is recreation, urban protection, some urban use, and some forestry.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 0%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (39 mm)

**Profile readily available water (0 – 100 cm):** Very low (8 mm)

**Topsoil bulk density:** 1.29 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.48 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 0.5 – 1.7%

**Topsoil P retention:** Very low (3%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Ohope sand** is the predominant soil type. Areas without topsoil especially in blow-outs also occur.

#### ***Associated and similar soils***

**Papamoa series** occur on more inland dunes with deeper topsoils and more organic matter; **Piripai series** and **Kopeopeo series** with thin layers of tephra.

## **General land use suitability ratings**

### **Ohope sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Low water-holding capacity, accelerated wind erosion risk, very low fertility.
Horticulture	Not suitable	Low water-holding capacity, accelerated wind erosion risk, very low fertility.
Intensive pasture	Not suitable	Low water-holding capacity, accelerated wind erosion risk, very low fertility.
Forestry	Not suitable to low	Low water-holding capacity, accelerated erosion risk, very low fertility.

### ***Management practices to improve suitability***

- Employ protection planting to combat wind erosion.



**Ohope sand (scale in m)**

## Soil Series Name: Okoia (Ok)

### Overview

Okoia soil series occur on flattish swales behind fore dunes in coastal areas in the Cape Runaway area. The soils are formed from alluvium and wind-blown sand. Soil profiles consist of dark brown mottled silt loam on grey mottled clay loam with abundant weathered rounded greywacke gravels below 60 cm. The soils are classified as **Typic Orthic Gley Soils**. Land use is rough pasture (dry stock) and some maize cropping.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 24%

**Potential rooting depth:** 50 – 70 cm

**Rooting barrier:** Fluctuating high ground water table

**Drainage class:** Poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Moderate to low (67 mm)

**Profile readily available water (0 – 100 cm):** Moderate (56 mm)

**Topsoil bulk density:** 0.91 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** Not known

### Soil types/variations

**Okoia silt loam** is the main soil type. Mottles in the topsoil occur in very wet places. Peaty layers also occur locally.

#### ***Associated and similar soils***

**Opoutama series** on coastal flats; **Oweka series** on adjacent alluvial flats.

## General land use suitability ratings

### Okoia silt loam

Land use	Suitability rating	Management considerations
Arable	Not suitable to low	Poor natural drainage, often difficult to drain, low fertility.
Horticulture	Not suitable	Poor natural drainage, often difficult to drain, low fertility.
Intensive pasture	Low	Poor natural drainage often difficult to drain.
Forestry	Not suitable	Poor natural drainage.

### Management practices to improve suitability

- Use zero or minimum tillage methods for fodder cropping.
- Employ drainage where economically and physically feasible.
- Use as good wetland preservation areas.



Okoia silt loam

## Soil Series Name: Omarumutu (Om)

### Overview

Omarumutu soil series occur on flattish swales behind fore dunes in coastal areas in the Opotiki area and on Matakana Island near Tauranga. The soils are formed from wind-blown sand with a thin cover of rhyolitic tephra (Kaharoa, Taupo and Whakatane Tephra). Soil profiles consist of very dark brown sandy loam on dark brown mottled loamy sand, on brown mottled coarse sand (Taupo Pumice) overlying greyish brown and olive sand. The soils are classified as **Typic Sandy Gley Soils**. Land use is rough pasture with rushes (dry stock) and some areas are in flax.

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 10 – 15%

**Potential rooting depth:** 40 – 60 cm

**Rooting barrier:** Fluctuating high ground water table

**Drainage class:** Poorly drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (105 mm)

**Profile readily available water (0 – 100 cm):** Moderate (36 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.38 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 2.6 – 7.8%

**Topsoil P retention:** Medium (42%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Omarumutu sandy loam** is the main soil type. In this series, the amount of tephra varies and in places, Kaharoa Tephra is missing, or there is less Taupo Tephra. Iron accumulation can also occur in the subsoil.

### *Associated and similar soils*

**Papamoa series** without mottles or tephra on subdued dunes; **Ohope series** on sandy fore dunes.

## General land use suitability ratings

### Omarumutu sandy loam

Land use	Suitability rating	Management considerations
Arable	Not suitable to low	Poor natural drainage, often difficult to drain, low fertility.
Horticulture	Not suitable	Poor natural drainage, often difficult to drain, low fertility.
Intensive pasture	Low	Poor natural drainage often difficult to drain.
Forestry	Not suitable	Poor natural drainage.

### Management practices to improve suitability

- Use zero or minimum tillage methods for fodder cropping.
- Employ drainage where economically and physically feasible.
- Use as good wetland preservation areas.



Omarumutu sandy loam (scale in m)

## Soil Series Name: Omeheu (Ome)

### Overview

Omeheu soil series occur on flat low-lying terraces in western parts of the Rangitaiki Plains. Parent materials are very thin Tarawera Tephra on sandy Kaharoa alluvium, on Kaharoa Tephra on pumice alluvium or peat. Soil profiles show deep very dark brown loamy sand, overlying pale olive fine loamy sand and light grey mottled sandy loam, resting on dark greyish brown sandy clay loam and light grey sand. The soils are classified as **Acidic Orthic Gley Soils**. The soils are used for dairying, beef, cropping (maize), and horticulture (kiwifruit, feijoas, boysenberries).

### Physical properties

**Texture:** Sand over loam

**Topsoil clay content:** 5 – 7%

**Potential rooting depth:** Unlimited to 1m.

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Imperfectly drained

**Permeability:** Rapid over moderate

**Profile total available water (0 – 100 cm):** High (232 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (88 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** High P; Low Ca and Mg; high K

### Soil types/variations

**Omeheu gritty loamy sand** is the main soil type. In this soil series, there is variation in the thickness of layers. For example, the Kaharoa Tephra-derived alluvium ('buff layer') varies from 20 to 50 cm in thickness.

#### ***Associated and similar soils***

**Awaiti series** occur at somewhat higher levels and have sandier soil textures. **Paroa series** occur in lower-lying depressions and have a silt loam to silty clay 'buff layer'.

## **General land use suitability ratings**

### **Omeheu gritty loamy sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Poor to imperfect natural drainage, low fertility, fragile topsoil.
Horticulture	Moderate to low	Poor to imperfect natural drainage, low fertility, fragile topsoil.
Intensive pasture	Moderate	Poor to imperfect natural drainage.
Forestry	Not suitable	Poor to imperfect natural drainage.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for cropping.
- Employ drainage where possible (open drains).
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Omeheu loamy sand**

## Soil Series Name: Onepū (Onp)

### Overview

Onepu soil series occur on river terraces on the Rangitaiki Plains in the Onepu and Te Teko localities. Parent materials are Tarawera Tephra on sandy and fine silty alluvium. Soil profiles show deep very dark brown sand with many fine Tarawera lapilli, overlying greyish-brown silt loam. They rest on light grey coarse sand and light brownish grey silt. Compact grey fine sand occurs below 80 cm depth. The soils are classified as **Acidic Orthic Gley Soils**. Land use is dairying, cropping (maize), horticulture (pip and stone fruit), carrots, tomatoes, strawberries, and onions.

### Physical properties

**Texture:** Sand over loam

**Topsoil clay content:** 5 – 8%

**Potential rooting depth:** About 70 – 100 cm

**Rooting barrier:** Fluctuating ground water table

**Drainage class:** Poorly to imperfectly drained

**Permeability:** Rapid over moderate

**Profile total available water (0 – 100 cm):** Moderate (117 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (77 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** Not known but possibly low

### Soil types/variations

**Onepu sand** is the main soil type. There is variation in the thickness of layers. Inclusions of mottled **Kawerau series** occur.

#### *Associated and similar soils*

**Maketu series** have diatomaceous earth layers and generally finer textures.

## **General land use suitability ratings**

### **Onepu sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Poor to imperfect natural drainage, low fertility, fragile topsoil.
Horticulture	Moderate	Poor to imperfect natural drainage, low fertility, fragile topsoil.
Intensive pasture	Moderate	Poor to imperfect natural drainage.
Forestry	Not suitable	Poor to imperfect natural drainage.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for cropping.
- Employ drainage where possible.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Onepu sand**

## Soil Series Name: Opotiki (Op)

### Overview

Opotiki soil series occur on flat to rolling and hilly land in the Opotiki region. They are formed from very thin Taupo Tephra on rhyolitic tephra and loess. Soil profiles show black sandy loam topsoils on dark yellowish-brown sandy loam resting on yellowish-brown sandy loam and silt loam. The soils are classified as **Allophanic Orthic Pumice Soils**. Land use includes kiwifruit orchards, some citrus orchards, dairying and dry stock.

### Physical properties

**Texture:** Loam over sand and loam

**Topsoil clay content:** 8 – 15%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid (moderate on hilly slopes)

**Profile total available water (0 – 100 cm):** High (152 mm)

**Profile readily available water (0 – 100 cm):** High (54 – 64 mm)

**Topsoil bulk density:** 0.91 g/cm<sup>3</sup>

**Subsoil bulk density:** 0.84 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** High (57 – 73%)

**Available P, Ca, Mg and K:** Low (strongly leached soil)

### Soil types/variations

**Opotiki sandy loam** occurs on flat to undulating surfaces, **Opotiki sandy loam, rolling phase** occurs on rolling land, and **Opotiki hill soils** are on hilly slopes. Thinner tephra layers on hilly slopes occur which have clay loam texture in lower subsoils. Rolling phase soils can have silt loam topsoils.

### *Associated and similar soils*

**Whakatane series** have coarser textures and a wider climatic range. **Te Puke series** are derived from different rhyolitic tephra.

## **General land use suitability ratings**

### **Opotiki sandy loam and Opotiki sandy loam, rolling phase**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Fragile topsoil, low fertility.
Horticulture	High	Low fertility.
Intensive pasture	High	Low fertility.
Forestry	High	No limitations.

### **Suitability rating Opotiki hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Hilly slopes, fragile topsoil, low fertility.
Horticulture	Low	Hilly slopes, fragile topsoil, low fertility.
Intensive pasture	High	Hilly slopes, low fertility.
Forestry	High	No limitations.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Opotiki sandy loam**

## Soil Series Name: Opouriao (Ou)

### Overview

Opouriao soil series occur on levees of river terraces in the Opotiki area, the Rangitaiki Plains and Cape Runaway area. The soils are derived from alluvium from greywacke and tephra. Common soil profiles have 23 cm black fine sandy loam or silt loam which rest on yellowish-brown to olive brown fine sandy loam overlying yellowish-brown silt loam. The soils are classified as **Weathered Fluvial Recent Soils** and belong to the most versatile soils in New Zealand (similar, for example, to **Manawatu soils**). All soils would flood infrequently without stop bank protection. Land use is dairying, cropping (maize), and horticulture (kiwifruit, carrots, tomatoes, strawberries, onions).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 10 – 24%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** High (209 mm)

**Profile readily available water (0 – 100 cm):** Very high (57 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 8.6%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** High P; Medium to high Ca, Mg and K

### Soil types/variations

**Opouriao fine sandy loam** and **silt loam** have been mapped. **Opouriao mottled fine sandy loam** has fine strong brown mottles below 40 cm. Locally, there are rounded greywacke gravels at the surface especially near the edge of the levee. A gravelly variant was mapped (in detailed surveys) on stony ridges.

### *Associated and similar soils*

**Rangitaiki series** occur on lower river terraces, usually adjacent to the river. These would flood frequently without stop bank protection. **Otara series** are moderately gleyed and occur adjacent but further inland of Opouriao series. **Ruatoki series** occur below Opouriao series on a lower river terrace.

## ***General land use suitability ratings***

### **Opouriao fine sandy loam and silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	High	Fragile topsoil.
Horticulture	High	Fragile topsoil, slight flooding risk.
Intensive pasture	High	No limitations.
Forestry	Moderate to high	Slight flooding risk.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Opouriao fine sandy loam (scale in m)**

## Soil Series Name: Opoutama (Ox)

### Overview

Opoutama soil series occur in areas south of Cape Runaway and in Waihou Bay on low-angle dunes. Parent material is wind-blown sand. Soil profiles have thin (5 cm) dark brown coarse sand resting on olive brown and light yellowish-brown coarse sand. The soils are classified as **Sandy Raw Soils**. Land use is recreation. Some areas are in rough pasture.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 0 – 2%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (40 mm)

**Profile readily available water (0 – 100 cm):** Low to very low (8 – 12 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 0.5 – 1.75%

**Topsoil P retention:** Very low (3%)

**Available P, Ca, Mg and K:** Not known, but probably low

### Soil types/variations

**Opoutama sand** is the main soil type. Small areas with thicker topsoil (17 cm black peaty loamy sand overlying dark greyish brown sand) were mapped as **Opoutama black sand** and classified as **Typic Sandy Brown Soils**.

#### ***Associated and similar soils***

**Ohope series** near Whakatane and Tauranga; **Okoia silt loam** on shallow swales behind the dunes.

## **General land use suitability ratings**

### **Opoutama sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low to not suitable	Erosion acceleration, low water-holding capacity; low fertility, fragile topsoil, saline winds.
Horticulture	Not suitable	Erosion acceleration, low water-holding capacity, low fertility, fragile topsoil, saline winds.
Intensive pasture	Low	Low water-holding capacity, low fertility.
Forestry	Low	Low water-holding capacity, saline winds.

### ***Management practices to improve suitability***

- Employ protection vegetation to prevent wind erosion.



**Opoutama sand**

## Soil Series Name: Orangihewa (Og)

### Overview

Orangihewa soil series occur on flat terraces in the southern parts of the Opotiki flats. They are formed from very thin Taupo Tephra on water-sorted Whakatane Tephra overlying alluvium derived from greywacke and tephra, resting on greywacke gravels. Profiles have thick (26 cm) black loamy sand topsoils overlying dark yellowish-brown loamy sand and yellowish-brown firm loamy sand. The soils are classified as **Allophanic Orthic Pumice Soils**. Land use is dry stock and some cropping (maize).

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 5 – 10%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (96 mm)

**Profile readily available water (0 – 100 cm):** Moderate (72 mm)

**Topsoil bulk density:** 0.91 g/cm<sup>3</sup>

**Subsoil bulk density:** 0.84 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** High (65%)

**Available P, Ca, Mg and K:** Medium P; low Ca, Mg and K (strongly leached soil)

### Soil types/variations

**Orangihewa loamy sand** is the main soil type. Inclusions of **sandy loam** and **fine sandy loam** topsoils occur.

#### ***Associated and similar soils***

**Opouriao series** are younger soils on lower river terraces.

## **General land use suitability ratings**

### **Orangihewa loamy sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Fragile topsoil, low fertility.
Horticulture	Not suitable to low	Fragile topsoil, low fertility.
Intensive pasture	Moderate to high	Low fertility.
Forestry	High	No limitations.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Orangihewa loamy sand**

## Soil Series Name: Oruanui (Oi)

### Overview

Oruanui soil series occur on flat, rolling and hilly land in southern parts of the Bay of Plenty. The soils are formed from more than 50 cm Taupo Pumice on older weathered rhyolitic tephra. The original vegetation was podocarp forest under 1500 to 1800 mm annual rainfall. This resulted in reddish-looking subsoil. Profiles show black to very dark grey sandy topsoils overlying dark brown and dark reddish-brown sand, which rest on strong brown to yellowish-brown loamy sand. The underlying weathered tephra is brown greasy silt loam. The soils are classified as **Podzolic Orthic Pumice Soils**. Land use is dry stock, fodder cropping, dairying and exotic forestry.

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 5 – 15%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to high (128 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (80 mm)

**Topsoil bulk density:** 0.91 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low (strongly leached soil)

### Soil types/variations

**Oruanui sand** occurs towards Taupo with deeper Taupo Pumice and including a thick lapilli layer. **Oruanui sandy loam** generally has a thinner layer of Taupo Pumice (but more than 50 cm) overlying weathered rhyolitic tephra. **Oruanui hill soils** are on hilly slopes with 50 to 210 cm Taupo Pumice overlying weathered rhyolitic tephra.

### *Associated and similar soils*

**Taupo series** occur under lower annual rainfall and shrub vegetation, resulting in thinner topsoils and B horizons (about 10 cm). **Tihoi series** occur under 1800 mm or more annual rainfall and a rimu vegetation resulting in Podzols.

## **General land use suitability ratings**

### **Oruanui sand and sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Fragile topsoil, low fertility, cool climate.
Horticulture	Low to not suitable	Fragile topsoil, low fertility, cool climate.
Intensive pasture	Moderate to low	Low fertility, cool climate.
Forestry	High	No limitations.

### **Oruanui hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, fragile topsoil, low fertility, cool climate.
Horticulture	Not suitable	Hilly slopes, fragile topsoil, low fertility, cool climate.
Intensive pasture	Low to moderate	Hilly slopes, low fertility, cool climate.
Forestry	High	Cool climate, hilly slopes.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Oruanui sand** (Note the deep black topsoil, red B horizons, and grey Rotongaio Ash at 90 cm, scale in m)

## Soil Series Name: Otara (Ota)

### Overview

Otara soil series occur on former back swamps of the flood plains of the Waioeka and Otara Rivers in the Opotiki area. The soils are derived from alluvium derived from greywacke and tephra. The soils have deep very dark greyish brown friable silt loam with moderately developed structure, resting on black silt loam with a few gravels on light olive brown mottled fine sandy loam on light olive brown to pale olive mottled silt loam. The soils are classified as **Mottled Fluvial Recent Soils**. Land use is dairying, dry stock, fodder cropping (maize) horticulture and market gardening (tomatoes, beans).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 19 – 22%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Imperfectly drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** High (175 mm)

**Profile readily available water (0 – 100 cm):** High (124 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 8.6%

**Topsoil P retention:** Medium (33%)

**Available P, Ca, Mg and K:** Medium (weakly leached soil)

### Soil types/variations

**Otara silt loam** is the main soil type. Inclusions of soils with clay loam and sandy subsoils are present.

#### ***Associated and similar soils***

**Opouriao series** occur on well drained levees (no mottling in the subsoil). **Apanui series** and **Waioeka series** are poorly drained and occur on former back swamps. Subsoil colours are greyer and mottles brighter.

## General land use suitability ratings

### Otara silt loam

Land use	Suitability rating	Management considerations
Arable	High	Imperfect drainage, locally greywacke gravels at the surface.
Horticulture	High	Imperfect drainage.
Intensive pasture	High	Imperfect drainage.
Forestry	Not suitable	Imperfect drainage.

### Management practices to improve suitability

- Use zero or minimum tillage methods for cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.
- Surface drainage.



Otara silt loam (scale in m)

## Soil Series Name: Oweka (Ow)

### Overview

Oweka soil series occur on flood plains of streams and river of Eastern Bay of Plenty with moderate areas south of Cape Runaway. The soils are derived from alluvium derived from greywacke and tephra. Soil profiles have dark brown friable silt loam resting on brown silt loam and fine sandy loam. Olive brown sand occurs at about 80 cm depth. The soils are classified as **Typic Fluvial Recent Soils**. Land use is dairying, dry stock, and fodder cropping (maize).

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** High (163 mm)

**Profile readily available water (0 – 100 cm):** High (101 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 8.6%

**Topsoil P retention:** Low (19%)

**Available P, Ca, Mg and K:** Medium (weakly leached soil)

### Soil types/variations

**Oweka silt loam** is the main soil type. Inclusions of **sandy loam** topsoils and rounded greywacke gravels occur.

#### ***Associated and similar soils***

**Waiapu series** on the lower flood plain of the river flats with sandier and stony profiles often without distinct topsoil; **Waihoata series** on poorly drained back swamps.

## General land use suitability ratings

### Oweka silt loam

Land use	Suitability rating	Management considerations
Arable	Moderate to high	Infrequent flooding risk, locally stones on the surface.
Horticulture	Moderate to low	Infrequent flooding risk, locally stones on the surface, cool climate.
Intensive pasture	Moderate	Infrequent flooding risk.
Forestry	Not suitable	Infrequent flooding risk.

### Management practices to improve suitability

- Use zero or minimum tillage methods for cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.



**Oweka silt loam** (Note rounded greywacke gravels forming part of the matrix of this profile, spade 90 cm long)

## Soil Series Name: Paerata (Pe)

### Overview

Paerata soil series occur on flat to hummocky valley floors in the Opotiki area. Parent materials are alluvium derived from rhyolitic tephra including very thin Kaharoa Tephra, on peat, on more alluvium with very thin Taupo Tephra on alluvium with wood remnants. Soil profiles are multilayered and briefly consist of dark brown peaty silt loam on light grey sand (Kaharoa Tephra) on dark reddish-brown peat. This overlies light olive grey silt loam mixed with Taupo lapilli, resting on pale yellow coarse sand (Taupo Tephra) on alluvium with wood remnants. The soils are classified as **Peaty Acid Gley Soils**. Land use is predominantly pasture (dairying, dry stock) with very small areas in crop (maize). Blueberry trials have been done.

### Physical properties

**Texture:** Loam over peat

**Topsoil clay content:** 20 – 30%

**Potential rooting depth:** 80 – 90 cm

**Rooting barrier:** Fluctuating ground water table

**Drainage class:** Poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** High (161 mm)

**Profile readily available water (0 – 100 cm):** High (103 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.38 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 10.3 – 13.8%

**Topsoil P retention:** Medium (52%) in peaty horizon, low to very low in other horizons

**Available P, Ca, Mg and K:** Medium (moderately to weakly leached soil)

### Soil types/variations

**Paerata peaty silt loam** is the main soil type. Variations in thickness and occurrence of the soil layers occur as inclusions.

#### *Associated and similar soils*

**Hanaia series** are gley soils without peaty layers. **Taho series** are gley soils derived from older (than Taupo Tephra) rhyolitic tephra. **Matao series** are organic soils of the wetter parts of the valley floors.

## General land use suitability ratings

### Paerata peaty silt loam

Land use	Suitability rating	Management considerations
Arable	Moderate	Poor natural drainage, infrequent flooding risk.
Horticulture	Moderate	Poor natural drainage, infrequent flooding risk.
Intensive pasture	Moderate	Poor natural drainage, infrequent flooding risk.
Forestry	Not suitable	Poor natural drainage.

### Management practices to improve suitability

- Employ surface drainage. Drains should cut through natural well drained horizons (Kaharoa and Taupo Tephra layers).



Paerata peaty silt loam (Note Kaharoa Tephra at 20 cm depth, scale in m)

## Soil Series Name: Paroa (Pr)

### Overview

Paroa soil series occur on former back swamps of the Rangitaiki Plains where they are extensive. The soils are formed from layers of tephra (thin Tarawera, Kaharoa and Taupo Tephra), pumiceous alluvium, silt, and peat. In the lower subsoil, pumiceous sand and gravel may occur. Soil profiles have 23 cm very dark brown silt loam on pale yellow silt loam. Often, this layer of alluvium derived from Kaharoa Tephra is compact and is called the 'buff layer'. It overlies light grey loose sand (Kaharoa Tephra) on dark brown massive peat. The soils are classified as **Acidic Recent Gley Soils**. Land use is dairying, cropping (maize) and some dry stock.

### Physical properties

**Texture:** Loam over peat

**Topsoil clay content:** 20 – 30%

**Potential rooting depth:** 20 – 50 cm

**Rooting barrier:** Fluctuating ground water table

**Drainage class:** Poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Moderate (120 mm)

**Profile readily available water (0 – 100 cm):** Low (47 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.0 – 13.8%

**Topsoil P retention:** Medium (35%)

**Available P, Ca, Mg and K:** Low to medium

### Soil types/variations

**Paroa silt loam** has a silt loam subsoil. **Paroa silt loam on peat** has sandy layers on peat (most extensive type). **Paroa peaty silt loam on peat** has a peaty silt loam topsoil mixed with Tarawera lapilli. **Paroa coarse sandy loam** has layered sandy loam and silt loam textures. **Paroa coarse sandy loam on peat** is similar to coarse sandy loam but with peat below Kaharoa Tephra. **Paroa coarse sandy loam on gravel** has coarse sand and gravel below 90 cm depth.

### *Associated and similar soils*

**Pongakawa series** do not have the 'buff layer'. **Awakaponga series** occur closer to the river and flood infrequently and have several recent flood layers.

## **General land use suitability ratings**

### **Paroa silt loam and Paroa silt loam on peat**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Poor natural drainage, fluctuating ground water levels, fragile topsoil structure.
Horticulture	Moderate	Poor natural drainage, fluctuating ground water levels, fragile topsoil structure.
Intensive pasture	Moderate to high	Poor natural drainage, fluctuating ground water levels.
Forestry	Not suitable	Poor natural drainage, fluctuating ground water levels.

### **Management practices to improve suitability**

- Artificial drainage will bring these soils to moderately well drained status.



**Paroa silt loam on peat** (Note pale layer of Kaharoa Tephra, scale in inches)

## Soil Series Name: Piripai (Pi)

### Overview

Piripai soil series occur in a dune system with easy rolling slopes with ridges and swales along the coast of the Rangitaiki Plains. They are derived from wind-blown coastal sand from rhyolitic pumice alluvium, capped with a very thin (less than 30 cm) cover of basaltic Tarawera Tephra and rhyolitic Kaharoa Tephra. Soil profiles have thin black loamy sand on very dark grey coarse sand, resting on black loamy sand and dark brown sand. Subsoils are yellowish-brown sand. The soils are classified as **Sandy Raw Soils**. Land use is extensive grazing of dry stock and some dairying.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 4 – 10%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (98 mm)

**Profile readily available water (0 – 100 cm):** Moderate (63 mm)

**Topsoil available water (0 – 30 cm):** Low (21 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 0.5 – 1.7%

**Topsoil P retention:** Very low (3%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

Piripai loamy fine sand is the main soil type.

#### ***Associated and similar soils***

Where the tephra cover eroded off (mostly fore dunes), the soils are **Pikowai series**. Piripai soils are also associated with unnamed soils in swales where topsoils can be 30 cm thick.

## General land use suitability ratings

### Piripai loamy fine sand

Land use	Suitability rating	Management considerations
Arable	Low	Fragile topsoil structure, susceptible to severe wind erosion when cultivated, summer droughts, low fertility.
Horticulture	Not suitable	Fragile topsoil structure, susceptible to severe wind erosion when cultivated, summer droughts, low fertility.
Intensive pasture	Low to moderate	Summer droughts, low fertility.
Forestry	Moderate	Low fertility, fragile topsoil, wind erosion potential.

### Management practices to improve suitability

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.
- Forestry: avoid windrowing and damage to topsoils.



Piripai loamy fine sand

## Soil Series Name: Pongakawa (Po)

### Overview

Pongakawa soil series occur on the Rangitaiki Plains in Whakatane West, Awakeri, Paroa, Thornton and Greig Road localities. The soils are formed from fibrous sedge peat admixed with Tarawera and Kaharoa Tephra with small amounts of fine alluvium and a thin layer of Taupo Tephra. Profiles have black peaty sand on pale brown sand (Kaharoa Tephra) resting on dark reddish brown firm peat. The soils are classified as **Acid Humic Organic Soils**. Land use consists of dairying, dry stock, cropping (maize) and some market gardening (potatoes, tomatoes).

### Physical properties

**Texture:** Peat over sand

**Topsoil clay content:** 1 – 3%

**Potential rooting depth:** 30 – 40 cm

**Rooting barrier:** Fluctuating ground water table

**Drainage class:** Poorly drained

**Permeability:** Moderate.

**Profile total available water (0 – 100 cm):** High (161 mm)

**Profile readily available water (0 – 100 cm):** Moderate (67 mm)

**Topsoil bulk density:** 0.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 0.18 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 34.5 - 69%

**Topsoil P retention:** High (62%)

**Available P, Ca, Mg and K:** Low P and Ca; high Mg and K

### Soil types/variations

**Pongakawa peaty sand** is the main soil type. The peat deposits range from 3 to 6 m in thickness and the thin alluvial beds are in the upper 25 cm.

#### ***Associated and similar soils***

**Kopeopeo series** are located on adjacent dunes. **Paroa series** occur on slightly higher elevations with alluvial layers in the profile.

## General land use suitability ratings

### Pongakawa peaty sand

Land use	Suitability rating	Management considerations
Arable	Low to moderate	Poor natural drainage, fluctuating ground water levels, fragile topsoil structure, wet soil in spring makes it a late soil for cropping.
Horticulture	Moderate to low	Poor natural drainage, fluctuating ground water levels, fragile topsoil structure.
Intensive pasture	Moderate to high	Poor natural drainage, fluctuating ground water levels, summer droughts.
Forestry	Not suitable	Poor natural drainage, fluctuating ground water levels.

### Management practices to improve suitability

- Employ artificial drainage.



Pongakawa peaty sand

## Soil Series Name: Poronui (Poi)

### Overview

Poronui soil series occur along streams and rivers on river terraces no longer subject to flooding in the southern and eastern parts of the Whakatane District. The soils are derived from water-sorted Taupo Tephra. Profiles have very dark brown sand overlying dark yellowish brown sand on yellowish-brown, pale yellow and dark grey coarse sand. The soils are classified as **Podzolic Orthic Pumice Soils**. Land use is forestry and some extensive dry stock or dairying with many areas in manuka shrub.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 0 – 5%

**Potential rooting depth:** Unlimited.

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (89 mm)

**Profile readily available water (0 – 100 cm):** Moderate (68 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Poronui sand** is the main soil type. Some soils have Kaharoa Tephra in the topsoil near Te Waiti.

#### ***Associated and similar soils***

**Rangitaiki series** occur on lower terraces subject to frequent flooding.

## **General land use suitability ratings**

### **Poronui sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low to unsuitable	Cold climate, fragile topsoil structure, low natural fertility.
Horticulture	Not suitable	Cold climate, fragile topsoil structure, low natural fertility.
Intensive pasture	Low	Cold climate with short growing season, low fertility.
Forestry	Moderate	Cold climate, low fertility.

### **Management practices to improve suitability**

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Plant green crops to increase organic matter and thicken topsoils.
- Forestry: avoid windrowing and damage to topsoils.



**Poronui sand** (Layering from flooding is accentuated by iron deposits)

## Soil Series Name: Pōtikirua (PotS)

### Overview

Potikirua soil series occur east of Cape Runaway on steep and very steep slopes with basalt as parent material. Soil profiles have dark brown silt loam overlying basaltic rock at 36 cm depth. The soils are classified as **Typic Orthic Recent Soils**. Land use is dry stock and many areas are in indigenous forest and shrub.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** 35 – 40 cm

**Rooting barrier:** Weathered rock

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (70 mm)

**Profile readily available water (0 – 100 cm):** Low (46 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 6.0%

**Topsoil P retention:** Low (22%)

**Available P, Ca, Mg and K:** High

### Soil types/variations

**Potikirua steepland soil** is the main soil type. Thin **Potikirua series** are on areas of exposed rock especially on very steep cliffs.

#### ***Associated and similar soils***

**Matakaoa series** are on rolling and hill country with a layer of tephra overlying basalt.

## **General land use suitability ratings**

### **Potikirua steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep and very slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low	Steep and very steep slopes, severe erosion potential.
Forestry	Low	Steep and very steep slopes, severe erosion potential, shallow soil profiles.

### **Management practices to improve suitability**

- Avoid over stocking.
- Use aerial methods of forest harvesting.



**Potikirua steepland soil landscape** (Note terracettes – an indication of creep erosion)

## Soil Series Name: Pukemaku (PkS)

### Overview

Pukemaku soil series occur in the western parts of Whakatane District, east and northeast of Mount Tarawera on steep and very steep valley and gully sides. The soils are formed from thin (up to 20 cm) Tarawera Tephra on Kaharoa and Taupo Tephra, on weathered rhyolitic tephra on ignimbrite. Profiles have very dark brown gritty sand on dark brown sand and fine gravel resting on brown greasy loamy sand and yellowish-brown to strong brown coarse sand. The soils are classified as **Typic Orthic Pumice Soils**. Land use is sheep farming and protection forestry (rimu–tawa).

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 0 – 5%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (85 mm)

**Profile readily available water (0 – 100 cm):** Moderate (63 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Pukemaku stepland soils** vary in total thickness of tephra overlying ignimbrite.

#### ***Associated and similar soils***

**Manawahe series** on rolling to hilly land, where less than 20 cm fine Tarawera Tephra overlies Kaharoa Tephra on weathered rhyolitic tephra; **Matahina series** on easy rolling to hilly slopes where less than 20 cm coarse Tarawera lapilli overlies Kaharoa Tephra and weathered rhyolitic tephra; **Whakatane series** on easy rolling to hilly slopes where very thin Tarawera Tephra and Kaharoa Tephra overlie weathered rhyolitic tephra; **Haroharo series** on steep and very steep slopes where thicker and coarser Tarawera and Kaharoa Tephra occur.

## **General land use suitability ratings**

### **Pukemaku steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep and very steep slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low	Steep and very steep slopes, severe erosion potential, low fertility.
Forestry	Low	Steep and very steep slopes, severe erosion potential.

### **Management practices to improve suitability**

- Avoid over stocking.
- Use aerial methods of forest harvesting.



**Pukemaku steepland soil landscape**

## Soil Series Name: Pukerimu (Px)

### Overview

Pukerimu soil series occur on easy rolling, rolling and hilly uplands in southern parts of the Whakatane District. The soils are formed from Taupo Tephra on Waimihia Tephra on older weathered rhyolitic tephra. They typically have pumice gravel (Taupo lapilli) at the surface (hill soils) or immediately below the topsoil. Soil profiles have dark brown loamy sand topsoils resting on dark reddish-brown loamy sand with many Taupo lapilli. These sit on top of strong brown pumice gravel on brownish-yellow pumice gravel. The soils are classified as **Podzolic Orthic Pumice Soils**. Land use is forestry, both indigenous (beech, rimu) and plantation (*Pinus radiata*) types.

### Physical properties

**Texture:** Sand and sand over skeletal (hill soils)

**Topsoil clay content:** 0 – 8%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (45 mm)

**Profile readily available water (0 – 100 cm):** Low (37 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 0.84 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low (strongly leached soils)

### Soil types/variations

**Pukerimu loamy sand** and **sandy loam** soil types occur. Bs horizons vary in colour from reddish brown to strong brown. **Pukerimu hill soils** occur on hilly landscapes.

### Associated and similar soils

**Tihoi series** have pumice lapilli further down the soil profile. **Oruanui series** occur at lower elevations with lower annual rainfall.

## **General land use suitability ratings**

### **Pukerimu loamy sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low to unsuitable	Cold climate, fragile topsoil structure, low natural fertility.
Horticulture	Not suitable	Cold climate, fragile topsoil structure, low natural fertility.
Intensive pasture	Low	Cold climate with short growing season, low fertility.
Forestry	Moderate	Cold climate, low fertility.

### **Pukerimu hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, cold climate, fragile topsoil structure, strong leaching.
Horticulture	Not suitable	Hilly slopes, cold climate, fragile topsoil structure, low natural fertility.
Intensive pasture	Low	Cold climate with short growing season, low fertility, hilly slopes.
Forestry	Moderate	Cold climate, low fertility, erosion prone.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Forestry: avoid windrowing and damage to topsoils.



**Pukerimu loamy sand**

## Soil Series Name: Pukeroa (Pu)

### Overview

Pukeroa soil series occur east of Te Puke on flat to gently sloping gully floors eroded out of tephra-covered terraces. Parent materials are very thin rhyolitic colluvium and tephra (Kaharoa and Taupo Tephra) on weathered rhyolitic tephra. Soil profiles have (18 cm) black sandy loam topsoils overlying dark yellowish-brown and pale brown loamy sand which rest on very pale brown sand and pale brown and white coarse sand (Kaharoa Tephra). These in turn rest on dark yellowish-brown sandy loam. The soils are classified as **Typic Orthic Pumice Soils**. Land use includes dairying, intensive dry stock, cropping (maize) and kiwifruit orchards.

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 10 – 18%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to high (142 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (98 mm)

**Topsoil bulk density:** 0.91 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Medium to low

### Soil types/variations

**Pukeroa sandy loam** is the main soil type. Pukeroa sandy loam, mottled phase has an olive yellow coloured subsoil with mottles.

#### ***Associated and similar soils***

**Parawhenuamea series** have thinner upper horizons within Kaharoa Tephra. **Ohinepanea series** occur on flat tephra-covered surfaces and are formed from Kaharoa Tephra.

## ***General land use suitability ratings***

### **Pukeroa sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Low nutrient levels, fragile topsoil, summer droughts.
Horticulture	Low	Low nutrient levels, fragile topsoil, summer droughts.
Intensive pasture	Moderate	Low nutrient levels, fragile topsoil, summer droughts.
Forestry	Moderate	Low nutrient levels, fragile topsoil, small areas within intensively farmed area.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.



**Pukeroa sandy loam** (The white band is Kaharoa Tephra)

## Soil Series Name: Pureora (Pur)

### Overview

Pureora soil series occur in the southern parts of Whakatane District on easy rolling and rolling uplands (greater than 700 m above sea level with over 2200 mm annual rainfall). Parent materials are Taupo Tephra overlying weathered rhyolitic and andesitic tephra on rhyolite or greywacke. Soil profiles show dark reddish-brown humic silt loam on greyish-brown loamy sand which rest on dark reddish-brown sandy loam on light yellowish-brown gravelly sand (Taupo lapilli), overlying dark greyish-brown sandy loam. The soils are classified as **Humose Orthic Podzols**. The soils are not used for agriculture because of cold climate and poor access. They are utilised for protection forestry, reserves, and recreation (e.g. hunting).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (88 mm)

**Profile readily available water (0 – 100 cm):** Moderate (61 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 15.5%

**Topsoil P retention:** Medium (42%)

**Available P, Ca, Mg and K:** Low (very strongly leached soil)

### Soil types/variations

**Pureora silt loam** is the main soil type. Peaty topsoils occur locally. The total thickness of Taupo Tephra varies from 54 cm to more than 100 cm.

#### *Associated and similar soils*

**Tihoi series** have similar soil profiles but occur at lower elevations (below 700 m). **Urewera series** are on steep and very steep slopes with thinner tephra overlying greywacke.


## **General land use suitability ratings**

### **Pureora silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Cold climate, inaccessibility, low nutrient levels, strong leaching, fragile topsoil.
Horticulture	Not suitable	Cold climate, inaccessibility, low nutrient levels, strong leaching, fragile topsoil.
Intensive pasture	Not suitable to low	Cold climate, inaccessibility, low nutrient levels, strong leaching, fragile topsoil.
Forestry	Low	Cold climate, inaccessibility, low nutrient levels, strong leaching, fragile topsoil.

### **Management practices to improve suitability**

- These areas are best left in protection forestry.



**Pureora silt loam**

## Soil Series Name: Rangitāiki (Ran)

### Overview

Rangitāiki soil series occur on flat terraces along streams and rivers throughout the Bay of Plenty. They flood frequently without stop bank protection. Parent material is alluvium derived from greywacke and tephra. Profiles show dark greyish-brown sand on dark brown and olive sandy loam, resting on light olive brown sand. The soils are classified as **Typic Fluvial Recent Soils**. Land use (which depends a little on soil texture) consists of dry stock or rough pasture and maize cropping.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 2 – 6%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (103 mm)

**Profile readily available water (0 – 100 cm):** Moderate (67 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 8.6%

**Topsoil P retention:** Low (19%)

**Available P, Ca, Mg and K:** Medium to low

### Soil types/variations

Soil types include **gravel, gravelly sand, sand, sandy loam** and **loamy sand**. The finer textures (sandy loam and loamy sand) are more suited to maize cropping.

### *Associated and similar soils*

**Ruatoki series** (in Opotiki) occur on slightly higher river terraces and have generally finer textures. **Opouriao series** (Opotiki area, East Coast, Rangitāiki plains) occur on levees and naturally flood infrequently. They have deep topsoils and are well developed soils.

**Awakaponga series** (Rangitāiki Plains) are poorly drained former back swamps on the same terrace level. **Oweka series** (East Coast) occur on slightly elevated terraces with deeper topsoils.

## **General land use suitability ratings**

### **Rangitaiki sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low to unsuitable	Coarse soil textures, stony in places, flooding without stop banks.
Horticulture	Not suitable	Coarse soil textures, stony in places, flooding without stop banks.
Intensive pasture	Low	Coarse soil textures, stony in places, flooding without stop banks.
Forestry	Not suitable	Coarse soil textures, stony in places, flooding without stop banks.

### **Management practices to improve suitability**

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.



**Rangitaiki sand**

## Soil Series Name: Raukūmara (RauS)

### Overview

Raukumara soil series occur in the East Coast area, east of State Highway 35 on steep and very steep slopes under 1400 to 2800 mm annual rainfall. Parent material is greywacke. Soil profiles have thin dark greyish-brown gravelly silt loam overlying olive brown stony silt loam. The soils are classified as **Typic Orthic Recent Soils**. The soils are mostly covered in bush and indigenous forest. Small areas are in pasture (sheep farming).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** Unlimited

**Rooting barrier:** Fractured rock

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (56 mm)

**Profile readily available water (0 – 100 cm):** Low (29 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 6.0%

**Topsoil P retention:** Low (22%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Raukumara steepland soils** are variable. Areas of exposed rock are common, especially on very steep cliffs. Thin tephra occurs on stable crests and spurs.

#### ***Associated and similar soils***

**Urewera series** occur above 600 m elevation. **Motu series** occur on hilly slopes with or without or with very patchy tephra on stable slopes. **Te Kaha series** have a variable layer of tephra.

## **General land use suitability ratings**

### **Raukumara steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep and very steep slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low	Steep and very steep slopes, moderate to severe erosion potential, low fertility.
Forestry	Low	Steep and very steep slopes, moderate to severe erosion potential.

### ***Management practices to improve suitability***

- Avoid over stocking.
- Use aerial methods of forest harvesting.
- Protection forests on very steep slopes.



**Raukumara steepland soil**

## Soil Series Name: Rewatu (Re)

### Overview

Rewatu soil series occur in flat low-lying parts of the eastern parts of the Rangitaiki Plains and in flat former back swamps of the valleys south and east of the Rangitaiki plains. Parent materials are alluvium and colluvium with a thin layer of Kaharoa Tephra. Soil profiles have black to very dark grey friable fine sandy loam on pale olive mottled firm silt loam with a 10 cm thick light grey sandy layer (Kaharoa Tephra). The lower subsoil is pale olive firm mottled clay loam. The soils are classified as **Typic Orthic Gley Soils**. Land use consists of dairying, dry stock. Maize cropping if artificially drained.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 15 – 20%

**Potential rooting depth:** 50 – 60 cm

**Rooting barrier:** Anoxic conditions

**Drainage class:** Poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Moderate to low (85 mm)

**Profile readily available water (0 – 100 cm):** Moderate (52 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** Medium levels of Ca and Mg., high K, low P

### Soil types/variations

**Rewatu fine sandy loam** is the main soil type. Variations in topsoil texture include silt loam, sandy loam and loamy sand. Mottling occurs in the topsoil.

#### ***Associated and similar soils:***

Opouriao series on adjacent well drained levees.

## **General land use suitability ratings**

### **Rewatu fine sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Poor natural drainage, weakly developed topsoil structure.
Horticulture	Low	Poor natural drainage, weakly developed topsoil structure.
Intensive pasture	Moderate	Poor natural drainage, weakly developed topsoil structure.
Forestry	Not suitable	Poor natural drainage.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Provide drainage (open drains).



**Rewatu fine sandy loam**

## Soil Series Name: Ruakituri (Ru)

### Overview

Ruakituri soil series occur on easy rolling, rolling and hilly upland at 450 to 670 m elevation under 1500 to 2200 mm annual rainfall in Urewera National Park, in south-eastern parts of Whakatane District. The soils are formed from thin (10 – 14 cm) Kaharoa Tephra on thin Taupo Tephra overlying weathered rhyolitic tephra. Profiles consist of light grey sand and yellowish-brown sand (Kaharoa Tephra) on dark reddish-brown greasy loamy sand. These overlie strong brown and yellowish-brown pumice gravel (Taupo lapilli) overlying yellowish-brown and olive brown sand or coarse sand. The soils are classified as **Podzolic Orthic Pumice Soils**. Vegetation includes rimu, totara, beech forest, and some cut-over bush.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 1 – 10%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low to moderate (52 – 78 mm)

**Profile readily available water (0 – 100 cm):** Low to moderate (37 – 57 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low (strongly leached soil)

### Soil types/variations

**Ruakituri sand** occurs on rolling land while **Ruakituri hill soils** are on hilly slopes. Kaharoa Tephra and Taupo Tephra are often thicker towards the foot of a slope.

#### ***Associated and similar soils***

**Pekepeke series** occur at lower rainfall and elevations. Their subsoils are paler and show less evidence of iron and humus accumulation in the subsoil. **Urewera series** are on steep and very steep slopes with thinner tephra overlying greywacke. **Matawai series** are even more podzolised and have less Kaharoa Tephra.

## **General land use suitability ratings**

### **Ruakituri sand**


<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Cold climate, low fertility, difficult access.
Horticulture	Not suitable	Cold climate, low fertility, difficult access.
Intensive pasture	Low	Cold climate, low fertility, difficult access.
Forestry	Moderate	Cold climate, low fertility, difficult access.

### **Ruakituri hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, cold climate, low fertility, difficult access.
Horticulture	Not suitable	Hilly slopes, cold climate, low fertility, difficult access.
Intensive pasture	Low	Hilly slopes, cold climate, low fertility, difficult access.
Forestry	Moderate	Cold climate, low fertility, difficult access.

### ***Management practices to improve suitability***

- Forestry: avoid windrowing and damage to topsoils.
- Employ protection forestry.



**Ruakituri sand**

## Soil Series Name: Rūātoki (Rti)

### Overview

Ruatoki soil series occur on flat terraces along the Waioeka and Otara Rivers in the Opotiki area. The soils occur on flat river terraces just above Rangitaiki series and below **Opouriao series**. They have deep (18 cm), dark brown silt loam topsoils on dark brown to olive brown sandy loam with few rounded greywacke gravels which rest on olive brown loose sand with few greywacke gravels increasing with increasing depth. The soils are **Typic Fluvial Recent Soils**. Land use consists of dairying, some dry stock and maize cropping. A few areas are in kiwifruit.

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 10 – 15%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to high (125 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (83 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 8.6%

**Topsoil P retention:** Low (19%)

**Available P, Ca, Mg and K:** Medium

### Soil types/variations

Textures include topsoils that vary from **fine sandy loam** to **silt loam**.

#### ***Associated and similar soils***

**Rangitaiki series** occur on slightly lower river terraces and have generally coarser textures. **Opouriao series** (Opotiki area, East Coast, Rangitaiki plains) occur on levees and naturally flood infrequently. They have deep topsoils and are well developed soils. **Oweka series** (East Coast) are on slightly elevated terraces with deeper topsoils.

## **General land use suitability ratings**

### **Ruatoki silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Weakly developed topsoil structure with possible low organic matter, flooding risk without stop banks.
Horticulture	Moderate	Weakly developed topsoil structure with possible low organic matter, flooding risk without stop banks.
Intensive pasture	Moderate to high	Weakly developed topsoil structure with possible low organic matter, flooding risk without stop banks.
Forestry	Not suitable	Flooding risk without stop banks.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Rotate land use.



**Ruatoki silt loam**

## Soil Series Name: Tahoe (Ta)

### Overview

Taho soil series occur in the Opotiki hill country where they have been mapped on flat to undulating fans and slumped valley heads. They are formed from colluvium derived from rhyolitic tephra. Soil profiles have very dark greyish-brown silt loam overlying pale brown to light yellowish-brown silt loam. These overlie light brownish-grey firm heavy silt loam and light grey clay loam. The soils are classified as **Typic Orthic Gley Soils**. Land use is dairying, dry stock and some kiwifruit.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 12 – 15%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Imperfectly to poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Moderate to low (69 mm)

**Profile readily available water (0 – 100 cm):** Low (48 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Taho silt loam** is the main soil type but textures range from sandy loam to clay loam in the upper horizons and loamy sand to clay in the subsoils. This is a result of older weathered tephra layers being eroded to the surface in a slump situation.

### *Associated and similar soils*

**Hanaia series** are derived mainly from tephric alluvium on the valley floors. **Iwiroa series** are derived from tephra and greywacke with greywacke gravels in the profile.

## **General land use suitability ratings**

### **Taho silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Imperfect to poor natural drainage (difficult to drain), low fertility.
Horticulture	Moderate to low	Imperfect to poor natural drainage (difficult to drain), low fertility.
Intensive pasture	Moderate to high	Imperfect to poor natural drainage (difficult to drain), pugging in winter time, low fertility.
Forestry	Not suitable	Imperfect to poor natural drainage.

### **Management practices to improve suitability**

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Employ surface drainage.



**Taho soil landscape**

## Soil Series Name: Takamore (Tsl)

### Overview

Takamore soil series occur in the Te Rereaurira Stream area, south east of Cape Runaway, and in small areas along the coast on low-angle fans with up to 5 degree slopes. The soils are formed from colluvium derived from greywacke, mudstone and sandstone. Soil profiles have dark brown gravelly silt loam with many weakly weathered greywacke gravels, overlying dark yellowish-brown stony clay loam with many weakly weathered rounded greywacke gravels. These overlie light brownish-grey clay loam. The soils are classified as **Typic Orthic Recent Soils**. Land use is dry stock farming and limited cropping (fodder crops).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid over moderate

**Profile total available water (0 – 100 cm):** Moderate to high (147 mm)

**Profile readily available water (0 – 100 cm):** moderate to high (82 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 6.0%

**Topsoil P retention:** Low (22%)

**Available P, Ca, Mg and K:** High P; medium to high Ca, Mg and K

### Soil types/variations

**Takamore gravelly silt loam** is the main soil type. This soil can occur in imperfectly drained parts of a fan, usually the lower parts.

#### *Associated and similar soils*

**Iwiroa series** are derived from tephra and greywacke with greywacke gravels in the profile. **Te Kaha series** and **Tokata series** are on terraces and valley floors.

## **General land use suitability ratings**

### **Takamore gravelly silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Moderate	Stony soil profiles, summer droughts.
Horticulture	Low	Stony soil profiles, imperfectly drained parts of the fans, summer droughts.
Intensive pasture	Moderate to high	Stony soils, summer droughts.
Forestry	Low	Summer droughts, soils occur in small areas only.

### **Management practices to improve suitability**

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Surface drainage on the wetter parts of the fans.



**Takamore soil landscape**

## Soil Series Name: Tarawera (Tr)

### Overview

Tarawera soil series occur on easy rolling, rolling and hilly country chiefly in Tarawera Forest in central Whakatane District. The soils are derived from more than 20 cm coarse Tarawera Tephra overlying Kaharoa Tephra. Soil profiles have black to dark greyish brown coarse Tarawera lapilli overlying black sand and yellowish-brown sand. These rest on yellowish-brown and pale yellow pumice gravel (Kaharoa Tephra). The soils are classified as **Buried-pumice Tephric Recent Soils**. Land use is forestry and dry stock.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 0 – 4%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (39 mm)

**Profile readily available water (0 – 100 cm):** Low (27 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.38 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (32%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Tarawera gravel** occurs on flat to rolling land. Tarawera Tephra varies in thickness from 23 to 90 cm. **Tarawera hill soils** are on hill slopes and Tarawera Tephra varies from 23 to 50 cm in thickness with accumulation of eroded material on foot slopes.

### *Associated and similar soils*

**Haroharo series** are on steep and very steep slopes with thinner layers of tephra overlying ignimbrite.

## **General land use suitability ratings**

### **Tarawera gravel**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Cool climate, low fertility, severe erosion potential.
Horticulture	Not suitable	Cool climate, low fertility, severe erosion potential.
Intensive pasture	Low	Low fertility, erosion potential on slopes over 10 degrees.
Forestry	Moderate	Low fertility.

### **Tarawera hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, cool climate, low fertility.
Horticulture	Not suitable	Hilly slopes, cool climate, low fertility, severe erosion potential.
Intensive pasture	Low	Hilly slopes, cold climate, low fertility, severe erosion potential.
Forestry	Low	Low fertility, severe erosion potential.

### ***Management practices to improve suitability***

- Limit cultivation to slopes below 10 degrees.
- Use low stocking rates with farming.
- Build up organic matter by ploughing in annual crops.
- Maintain a good pasture sward.
- Avoid stock tracking on steep slopes.
- Forestry: avoid windrowing and damage to topsoils.
- Employ protection forestry.



**Tarawera gravel** (Note thick coarse Tarawera Tephra and buried topsoil of Kaharoa Tephra)

## Soil Series Name: Tauhara (TaS)

### Overview

Tauhara soil series occur on steep and very steep slopes in the south-western part of Kaingaroa Forest. Parent materials are shallow to moderately thick Taupo Tephra overlying older weathered rhyolitic tephra. Soil profiles have thin very dark greyish-brown loose gritty sand on dark yellowish-brown loose gritty sand. These rest on light yellowish-brown loose gravelly sand and pumice gravel. The soils are classified as **Immature Orthic Pumice Soils**. Land use consists of forestry or dry stock farming.

### Physical properties

**Texture:** Sand over skeletal

**Topsoil clay content:** 0 – 5%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (54 mm)

**Profile readily available water (0 – 100 cm):** Low (39 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Tauhara stepland soils** have a full cover of Taupo Tephra of varying thickness.

#### ***Associated and similar soils***

**Motumoa stepland soils** have thinner, patchy and finer Taupo Tephra that overlies older weathered rhyolitic tephra. **Taupo** and **Oruanui series** are on rolling and hilly slopes with thicker Taupo Tephra. **Waipahihi series** are derived from water-sorted Taupo Tephra and occur in valley the floors.

## **General land use suitability ratings**

### **Tauhara steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep slopes, erosion potential.
Horticulture	Not suitable	Steep slopes, erosion potential.
Intensive pasture	Low	Steep slopes, erosion potential, low fertility.
Forestry	Low to moderate	Steep slopes, erosion potential.

### **Management practices to improve suitability**

- Employ aerial harvesting in forestry to prevent severe erosion.
- Use low dry stock stocking rates to prevent further erosion.



**Tauhara steepland soil landscape**

## Soil Series Name: Taupō (Tp)

### Overview

Taupo soil series occur in the southern Bay of Plenty on flat to rolling surfaces and hilly slopes under relatively low annual rainfall (1000-1400 mm). They are derived from Taupo Pumice, developed under shrub-type vegetation and typically have thin (10-15 cm) A and Bw horizons. The sandy soils have low water-holding capacity and summer droughts are common. Natural soil fertility is low. The soils are classified as **Immature Orthic Pumice Soils**. The soils are currently used for forestry and grazing dry stock with occasional fodder cropping. Conversion from forestry to dairying is occurring in places.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 2 – 7% (sand and loamy sand); 10 – 15% (sandy loam)

**Potential rooting depth:** Unlimited

**Rooting barrier:** None

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (63 mm)

**Profile readily available water (0 – 100 cm):** Low (46 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6-12.0%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Taupo loamy sand** and **Taupo sandy loam** have finer textures and generally slightly better water-holding capacity than **Taupo sand**. **Taupo hill soils** occur on hilly slopes with thinner Taupo Pumice overlying older tephra.

#### ***Associated and similar soils***

**Oruanui series** which developed under podocarp forest, under higher annual rainfall (1400 - 1600 mm), have thicker A and B horizons. They are less droughty compared to Taupo series. **Tauhara series** and **Motumoa series** are on steep slopes with thinner Taupo Pumice overlying older tephra. **Waipahihi series** are derived from water-sorted Taupo Pumice and occur on valley floors. These are lapilli-rich, droughty soils. **Ngakuru series** are derived from thin Taupo Pumice overlying weathered tephra.

## General land use suitability ratings

### Taupo sand and Taupo loamy sand

Land use	Suitability rating	Management considerations
Arable	Low	Droughty soil, weak topsoil structure; low fertility, wind erosion if cultivated during dry spells, low fertility.
Horticulture	Not suitable to low	Droughty soil, weak topsoil structure; low fertility, wind erosion if cultivated during dry spells, low fertility.
Intensive pasture	Low to moderate	Summer droughts, low fertility.
Forestry	Moderate	Droughts affecting young trees, low fertility.

### Taupo sandy loam

Land use	Suitability rating	Management considerations
Arable	Moderate	Weak topsoil structure; low fertility, wind erosion if cultivated during dry spells, low fertility.
Horticulture	Not suitable to low	Droughty soil, weak topsoil structure; low fertility, wind erosion if cultivated during dry spells, low fertility.
Intensive pasture	Moderate	Summer droughts, low fertility.
Forestry	Moderate to high	Low fertility.

### Taupo hill soils

Land use	Suitability rating	Management considerations
Arable	Not suitable	Slopes too steep to cultivate.
Horticulture	Not suitable	Hilly slopes, cool climate.
Intensive pasture	Low	Hilly slopes, summer droughts, low fertility.
Forestry	Moderate	Hilly slopes, low fertility.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Green crops to increase organic matter and thicken topsoils.
- Forestry: avoid windrowing and damage to topsoils.



**Taupo sandy loam**

## Soil Series Name: Tawhia (TyS)

### Overview

Tawhia soil series occur extensively in north-eastern parts of Whakatane District on steep and very steep valley sides and hill country under 1300 to 1800 mm annual rainfall. The soils are formed from thin Tarawera Tephra, on Kaharoa Tephra, Taupo Tephra and older weathered rhyolitic tephra on greywacke. Profiles have dark brown sandy loam topsoils overlying dark yellowish-brown fine sandy loam or silt loam. These overlie light olive brown to yellowish-brown silt loam with many weathered greywacke fragments increasing with increasing depth. The soils are classified as **Buried-allophanic Orthic Pumice Soils**. Land use is dry stock farming.

### Physical properties

**Texture:** Sand

**Topsoil clay content:**

**Potential rooting depth:** 50 – 90 cm

**Rooting barrier:** Fractured rock

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (91 mm)

**Profile readily available water (0 – 100 cm):** Moderate (59 mm)

**Topsoil bulk density:** 0.91 g/cm<sup>3</sup>

**Subsoil bulk density:** 0.84 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Tawhia stepland soils** have thick tephra on stable slopes but very thin tephra on eroded areas.

#### ***Associated and similar soils***

**Whakatane series** occur on flat, rolling and hilly areas with thicker tephra layers.

## **General land use suitability ratings**

### **Tawhia steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep slopes.
Horticulture	Not suitable	Steep slopes.
Intensive pasture	Low	Steep slopes, low fertility, moderate scree and slip erosion.
Forestry	Low	Steep slopes, moderate erosion risk, low fertility.

### **Management practices to improve suitability**

- Use aerial harvesting with forestry to reduce topsoil damage.
- Employ protection forestry.
- Use low stocking rates with dry stock farming to avoid further erosion.



**Tawhia steepland soil landscape** (Note terracettes indicating creep erosion)

## Soil Series Name: Te Kaha (TeK)

### Overview

Te Kaha soil series occur on easy rolling, rolling and hilly areas along the coast from Cape Runaway to western Hawai Bay. They are formed from rhyolitic tephra. Profiles show deep (20 cm) black sandy loam on dark brown sandy loam overlying dark yellowish silt loam at about 35 cm depth. The soils are classified as **Typic Allophanic Brown Soils**. Land use includes dairying, dry stock farming, kiwifruit orchards, maize and fodder cropping.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 22 – 26%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** High (165 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (95 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 9.5 – 19.0%

**Topsoil P retention:** Medium (55%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Te Kaha sandy loam** is the main soil type but silt loam-textured soils occur. Greywacke or siltstone gravels may occur below 1 m depth. Slightly thinner tephras occur on hilly slopes.

#### *Associated and similar soils*

**Raukumara series** are on steep and very steep slopes derived from greywacke.

#### *General land use suitability ratings*

##### Te Kaha sandy loam

Land use	Suitability rating	Management considerations
Arable	High	Low fertility, saline winds.
Horticulture	High	Low fertility, saline winds.
Intensive pasture	High	Low fertility.
Forestry	High	Low fertility, saline winds.

## Te Kaha hill soils

Land use	Suitability rating	Management considerations
Arable	Not suitable	Hilly slopes, low fertility.
Horticulture	Not suitable	Hilly slopes, low fertility.
Intensive pasture	High	Hilly slopes, low fertility.
Forestry	High	Low fertility.

### *Management practices to improve suitability*

- Forestry: avoid windrowing and damage to topsoils.
- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Use shelterbelts for horticulture.



**Te Kaha sandy loam**

## Soil Series Name: Te Piki (Pik)

### Overview

Te Piki soil series occur in the Te Piki area south-east of Cape Runaway on flat valley floors. Parent materials are peat, alluvium and rhyolitic tephra (Taupo Tephra). Soil profiles have dark reddish-brown loamy peat on strong brown coarse sand (Taupo Tephra) which overlies dark reddish-brown peat. The soils are classified as **Mellow Humic Organic Soils**. Land use consists of dairying or dry stock.

### Physical properties

**Texture:** Peat

**Topsoil clay content:** 0 – 3%

**Potential rooting depth:** 30 – 50 cm

**Rooting barrier:** Fluctuating high ground water table

**Drainage class:** Poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Very high (332 mm)

**Profile readily available water (0 – 100 cm):** Very high (126 mm)

**Topsoil bulk density:** 0.10 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 34.5 – 69.0%

**Topsoil P retention:** High (62%)

**Available P, Ca, Mg and K:** Low to medium

### Soil types/variations

**Te Piki** peaty loam is the main soil type. Some profiles have silty layers in the subsoil, twigs, roots and stumps below 100 cm depth.

#### *Associated and similar soils*

**Tokata series** occur in other parts of the same valley.

### **General land use suitability ratings**

#### **Te Piki peaty loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Poor drainage, high groundwater table in winter.
Horticulture	Low	Poor drainage, high groundwater table in winter.
Intensive pasture	Moderate	Poor drainage, high groundwater table resulting in pugging.
Forestry	Not suitable	Poorly drained.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods for fodder cropping.
- Avoid cultivating the soil during dry spells to prevent loss of organic matter.
- Employ drainage, but over-drainage will bring stumps to the surface. Strict water control is desirable.



**Te Piki peaty loam**

## Soil Series Name: Te Rahu (TRa)

### Overview

Te Rahu soil series occur on the Rangitaiki Plains on subdued inland dunes. Parent materials are thin or very thin layers of rhyolitic tephra (Tarawera, Kaharoa, Taupo) and thick Whakatane Tephra overlying wind-blown sand. The total thickness of tephra is generally 90 cm. Soil profiles have black loamy sand very friable topsoils on very dark brown sandy loam and brown coarse sand. These rest on reddish-brown loamy sand. Below 90 cm depth light grey loose dune sand occurs. The soils are classified as **Immature Orthic Pumice Soils**. Land use consists of beef, dairying and sheep farming. Summer droughts occur.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 4 – 8%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (95 mm)

**Profile readily available water (0 – 100 cm):** Moderate (70 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low (Strongly leached soil)

### Soil types/variations

**Te Rahu loamy sand** is the main soil type. Some profiles have deep topsoils (up to 50 cm thick). Peaty layers occur locally above the Whakatane Tephra.

#### ***Associated and similar soils***

**Pongakawa** series occur in depressions associated with the dunes. **Kopeopeo** series occur on somewhat younger inland dunes. They are also formed from tephra on wind-blown sand but without Whakatane Tephra.

## **General land use suitability ratings**

### **Te Rahu loamy sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Weakly developed topsoil structure, low natural fertility, summer droughts.
Horticulture	Low	Weakly developed topsoil structure, low natural fertility, summer droughts.
Intensive pasture	Moderate	Weakly developed topsoil structure, low natural fertility, summer droughts.
Forestry	Not suitable	Areas too small.

### **Management practices to improve suitability**

- Use zero or minimum tillage methods for cropping.
- Avoid working the soil during dry spells to prevent loss of organic matter.



**Te Rahu loamy sand**

## Soil Series Name: Te Teki (TtS)

### Overview

Te Teki soil series occur on steep and very steep slopes, chiefly in central and southern Kaingaroa Forest. The soils are formed from thin Taupo Tephra and Waimihia Tephra on older rhyolitic tephra overlying ignimbrite. Soil profiles have dark reddish-brown shallow topsoils overlying olive grey sand on dark brown sand. These rest on yellowish-brown sand and dark reddish-brown hard sandy loam overlying dark yellowish-brown and yellowish-brown loamy sand. The soils are classified as **Humose Orthic Podzols**. Land use consists of forestry and/or recreation and conservation.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 1 – 7%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (93 mm)

**Profile readily available water (0 – 100 cm):** Moderate (67 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 15.5 %

**Topsoil P retention:** Medium (42%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Te Teki stepland soils** as described above. Shallow profiles overlying ignimbrite occur where the tephra has eroded off the slope.

#### ***Associated and similar soils***

**Tihoi series** and **Pukerimu series** are on easy rolling and hilly slopes with thicker tephra layers. **Urewera series** occur on steep and very steep slopes where tephra overlies greywacke.

## **General land use suitability ratings**

### **Te Teki steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep and very steep slopes, cool climate, low natural fertility.
Horticulture	Not suitable	Steep and very steep slopes, cool climate, low natural fertility.
Intensive pasture	Low	Steep and very steep slopes, cool climate, low natural fertility.
Forestry	Moderate	Steep and very steep slopes, cool climate, low natural fertility.

### **Management practices to improve suitability**

- Use aerial harvesting methods for forestry.



**Te Teki steepland soil landscape**

## Soil Series Name: Te Teko (Tto)

### Overview

Te Teko soil series occur around and south of Te Teko on the Rangitaiki Plains, on flat alluvial terraces. Parent materials are 10 to 20 cm Tarawera Tephra, on very thin Kaharoa Tephra on mixed pumice and greywacke alluvium. Soil profiles consist of dark brown firm sand on very dark grey friable sand which rest on brown to olive brown sandy loam overlying pale olive fine sandy loam. The soils are classified as **Typic Orthic Pumice Soils**. Land use consists of dairying, dry stock, fodder crops, horticulture (watermelon, strawberries) and cropping (maize).

### Physical properties

**Texture:** Sand over loam

**Topsoil clay content:** 2 – 8% (sand); 6 – 11% (sandy loam)

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to high (148 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (85 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1 %

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** High in the topsoil but low in the subsoil

### Soil type/variations

**Te Teko sand** and **Te Teko sandy loam** are the main soil types.

#### ***Associated and similar soils***

**Rangitaiki series** are on lower flood plain terraces with AC profiles and frequent flooding occurs if not protected by stop banks. **Kawerau series** have a cover of coarse Tarawera Tephra overlying coarse textured subsoil derived from pumice alluvium.

## General land use suitability ratings

### Te Teko sand and Te Teko sandy loam

Land use	Suitability rating	Management considerations
Arable	Moderate	Weakly developed topsoil structure, low natural fertility.
Horticulture	Moderate to low	Weakly developed topsoil structure, low natural fertility.
Intensive pasture	Moderate	Weakly developed structure, low natural fertility, summer droughts.
Forestry	Moderate to high	Moisture deficiency at seedling growth stage.

### Management practices to improve suitability

- Use zero or minimum tillage methods for cropping.
- Avoid working the soil during dry spells to prevent loss of organic matter.



**Te Teko sand** (Note deep black coarse Tarawera Tephra overlying buried topsoil of Kaharoa Tephra)

## Soil Series Name: Tihoi (Toi)

### Overview

Tihoi soil series occur in southern parts of Whakatane District on easy rolling to rolling and hilly uplands. Soil profiles consist of dark reddish-brown very friable sandy loam on greyish-brown friable loamy sand resting on dark brown sandy loam on light olive firm sand. Lower subsoils are pale yellow pumice gravel (Taupo lapilli). The soils are classified as **Humose Orthic Podzols**. Land use consists of forestry and dry stock.

### Physical properties

**Texture:** Loam over sand (sandy loam); sand over skeletal (hill soils)

**Topsoil clay content:** 10 – 15% (sandy loam); 2-8% (hill soils)

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (90 mm), Low (56 mm), hill soils

**Profile readily available water (0 – 100 cm):** Moderate (63 mm), Low (27 mm), hill soils

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 15.5 %

**Topsoil P retention:** Medium (42%)

**Available P, Ca, Mg and K:** Low (Strongly leached soil)

### Soil types/variations

**Tihoi sandy loam** occurs on flat to rolling slopes. **Tihoi hill soils** occur on hilly slopes generally with thinner tephra layers.

#### ***Associated and similar soils***

**Oruanui series** occur at lower elevations and lower annual rainfall. **Pukerimu series** are soils where Taupo lapilli layers are at the surface. **Te Teki series** occur on steep and very steep slopes where shallower tephra overlies ignimbrite. **Ruakituri series** are soils where shallow or very shallow Kaharoa Tephra overlies Taupo Pumice.

## **General land use suitability ratings**

### **Tihoi sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Weakly developed topsoil structure, low natural fertility, cool climate.
Horticulture	Not suitable	Cool climate, weakly developed topsoil structure, low natural fertility.
Intensive pasture	Low	Cool climate, low natural fertility, summer droughts.
Forestry	Moderate	Cool climate.

### **Tihoi hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Hilly slopes, weakly developed topsoil structure, low natural fertility, cool climate.
Horticulture	Not suitable	Hilly slopes, cool climate, weakly developed topsoil structure, low natural fertility.
Intensive pasture	Low	Cool climate, low natural fertility, hilly slopes, summer droughts.
Forestry	Moderate	Cool climate, hilly slopes.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods.
- Avoid working the soil during dry spells to prevent loss of organic matter.



**Tihoi sandy loam** (Note pale horizon below black topsoil and red iron staining in subsoil, scale in m)

## Soil Series Name: Tokata (To)

### Overview

Tokata soil series occur on flat alluvial terraces of valley floors and river back swamps southeast and south of Cape Runaway. Parent materials are alluvium and colluvium. Soil profiles consist of very dark brown clay loam overlying grey mottled clay loam merging to grey mottled sandy clay and grey mottled sand. The soils are classified as **Typic Orthic Gley Soils**. Land use consists of dairying, dry stock, fodder crops, and maize.

### Physical properties

**Texture:** Loam over clay

**Potential rooting depth:** Unlimited

**Topsoil clay content:** 25 – 30%

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Poorly drained

**Permeability:** Moderate

**Profile total available water (0 – 100 cm):** Moderate to high (134 mm)

**Profile readily available water (0 – 100 cm):** Moderate to high (88 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 15.5 %

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** High

### Soil types/variations

**Tokata clay loam** has clay loam texture overlying sandy deposits. Profiles occur without the sandy lower subsoil.

#### *Associated and similar soils*

**Te Piki series** occur in peaty depressions of the same terrace.

#### *General land use suitability ratings*

##### Tokata clay loam

Land use	Suitability rating	Management considerations
Arable	Moderate to low	Poor drainage.
Horticulture	Low	Poor drainage.
Intensive pasture	Moderate	Poor drainage, pugging.
Forestry	Not suitable	Poor drainage.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods.
- Ensure adequate artificial drainage.



**Tokata clay loam** (Note intense mottling in the subsoil)

## Soil Series Name: Torere (Tor)

### Overview

Torere soil series occur on stony beaches along the East Coast. Parent materials are ocean-deposited gravel and colluvial and alluvial silts. Soil profiles consist of dark brown sandy loam with profuse rounded greywacke gravels and stones on greywacke gravels with 10% sandy loam. The soils are classified as **Fluvial Raw Soils** (Torere gravel) and **Typic Fluvial Recent Soil** (Torere stony sandy loam). Land use consists of rough grazing and building sites on Torere stony sandy loam.

### Physical properties

**Texture:** Skeletal

**Topsoil clay content:** 0% (gravel); 10 – 15% (stony sandy loam)

**Potential rooting depth:** 20 – 40 cm

**Rooting barrier:** Extremely gravelly

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Very low (1 – 16 mm)

**Profile readily available water (0 – 100 cm):** Very low (1 – 16 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** Very low (<2% )

**Topsoil P retention:** Very low to low (3 - 19%)

**Available P, Ca, Mg and K:** Not known

### Soil types/variations

Profiles on the beaches with almost 100% gravels are called **Torere gravel**. Some soils further inland have stony sandy loam textures and are known as **Torere stony sandy loam**.

### General land use suitability ratings

#### Torere gravel

Land use	Suitability rating	Management considerations
Arable	Not suitable	Profuse gravesl, saline climate, flooding.
Horticulture	Not suitable	Profuse gravels, saline climate, flooding.
Intensive pasture	Not suitable	Profuse gravels, saline climate, flooding.
Forestry	Not suitable	Profuse gravels, saline climate, flooding.

### Torere stony sandy loam

Land use	Suitability rating	Management considerations
Arable	Not suitable	Profuse gravels, saline climate.
Horticulture	Not suitable	Profuse gravels, saline climate.
Intensive pasture	Low	Profuse gravels, saline climate.
Forestry	Not suitable	Profuse gravels, saline climate.

### *Management practices to improve suitability*

- None recommended.



**Torere gravel**

## Soil Series Name: Tuparoa (TupS)

### Overview

Tuparoa soil series occur on steep and very steep slopes northeast of Waihou Bay at the East Coast. Parent materials are mudstones. Soil profiles consist of dark brown friable silt loam overlying yellowish-brown to light olive brown friable silt loam, resting on brownish-yellow clay loam with mudstone gravels increasing with depth. The soils are classified as **Typic Orthic Recent Soils**. Land use consists of dry stock and forestry.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** 50 – 60 cm

**Rooting barrier:** Weathered rock

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (85 mm)

**Profile readily available water (0 – 100 cm):** Moderate (51 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 6.0 %

**Topsoil P retention:** Low (22%)

**Available P, Ca, Mg and K:** Not known, but possibly medium levels

### Soil types/variations

**Tuparoa steepland soils** can have shallower profiles occurring with more mudstone gravels throughout the profile.

#### *Associated and similar soils*

**Mangaomeko series** occur on associated hilly slopes.

#### *General land use suitability ratings*

##### Tuparoa steepland soils

Land use	Suitability rating	Management considerations
Arable	Not suitable	Steep and very steep slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low	Steep and very steep slopes.
Forestry	Low	Steep and very steep slopes.

### ***Management practices to improve suitability***

- Use aerial harvesting methods for forestry.
- Employ low stocking rates for dry stock.



**Tuparoa steepland soil landscape**

## Soil Series Name: Tutaetoko (TutS)

### Overview

Tutaetoko soil series occur on steep and very steep slopes in the Opotiki area. Parent materials are rhyolitic tephra overlying greywacke, sandstone and/or gravels. Soil profiles consist of very dark brown friable sandy loam overlying yellowish brown to dark yellowish-brown friable silt loam, resting on brownish yellow clay loam at about 60 cm. The soils are classified as **Typic Orthic Recent Soils**. Land use consists of dry stock, forestry, and recreation (hunting).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 10 – 15%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Medium

**Profile total available water (0 – 100 cm):** Moderate high (85 mm)

**Profile readily available water (0 – 100 cm):** Moderate (56 mm)

**Topsoil bulk density:** 0.78 g/cm<sup>3</sup>

**Subsoil bulk density:** 0.86 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 12.9 – 22.4 %

**Topsoil P retention:** High (83%)

**Available P, Ca, Mg and K:** Not known, but possibly low levels

### Soil types/variations

**Tutaetoko steepland soils** are found on steep and very steep slopes.

#### *Associated and similar soils*

**Opotiki series** occur on adjacent rolling and hilly land with thicker tephra. **Iwiroa series** occur on associated fans.

#### *General land use suitability ratings*

##### **Tutaetoko steepland soils**

Land use	Suitability rating	Management considerations
Arable	Not suitable	Steep and very steep slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low	Steep and very steep slopes.
Forestry	Low	Steep and very steep slopes.

### ***Management practices to improve suitability***

- Use aerial harvesting methods for forestry.
- Employ low stocking rates for dry stock.



**Tutaetoko stepland soil landscape**

## Soil Series Name: Urewera (US)

### Overview

Urewera soil series occur on steep and very steep slopes in and around Urewera National Park. Parent materials consist of rhyolitic tephra overlying greywacke. Soil profiles are variable but generally have dark reddish-brown greasy sandy loam on dark yellowish-brown sandy loam which rest on yellowish red fine pumice gravel on dark yellowish brown and yellowish brown silt loam. The soils are classified as **Podzolic Orthic Pumice Soils** and shallower profiles are classified as **Typic Orthic Recent Soils**. Land use consists of dry stock, plantation forestry and large areas in indigenous forest.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 1 – 5%

**Potential rooting depth:** Unlimited, but 40 – 50 cm only on recent soils

**Rooting barrier:** Weathered rock

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (84 mm)

**Profile readily available water (0 – 100 cm):** Moderate (61 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1 %

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low (strongly leached soils)

### Soil types/variations

**Urewera steepland soils** are on steep and very steep slopes. Shallow and stony profiles with little tephra also occur.

#### ***Associated and similar soils***

**Opotiki series** are on rolling and hilly land with thicker tephra and occur under generally lower annual rainfall. **Matawai series** occur on higher elevations and high annual rainfall. **Iwiroa series** occur on associated fans.

## ***General land use suitability ratings***

### **Urewera steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep and very steep slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low	Steep and very steep slopes, cool or cold climate.
Forestry	Low	Steep and very steep slopes, cold climate.

### ***Management practices to improve suitability***

- Use aerial harvesting methods for forestry.
- Employ low stocking rates for dry stock.



**Urewera steepland soil landscape**

## Soil Series Name: Waiapu (Wu)

### Overview

Waiapu soil series occur on low river flats of the East Coast area. Parent materials are alluvial greywacke gravels and sands. Soil profiles consist of very dark greyish-brown friable sand on dark greyish-brown loamy sand overlying dark greyish-brown gravelly sand, very firm with abundant weakly weathered rounded greywacke gravels. The soils are classified as **Typic Fluvial Recent Soils** (Waiapu sandy loam) or **Fluvial Raw Soils** (Waiapu stony sand). Land use consists of rough grazing.

### Physical properties

**Texture:** Sand

**Potential rooting depth:** 40 – 50 cm

**Rooting barrier:** Densely packed gravels

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (42 mm)

**Profile readily available water (0 – 100 cm):** Low (34 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.38 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 8.6%

**Topsoil clay content:** 3 – 6%

**Topsoil P retention:** Low (19%)

**Available P, Ca, Mg and K:** Medium to low

### Soil types/variations

**Waiapu sandy loam** has sandy material over gravels. **Waiapu stony sand** has greywacke gravel at the surface.

#### ***Associated and similar soils***

**Torere series** have profiles on the beaches with almost 100% gravels (Torere gravel) and some further inland with some sandy loam (Torere stony sandy loam).

## **General land use suitability ratings**

### **Waiapu sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Profuse gravels, saline climate, flooding.
Horticulture	Not suitable	Profuse gravels, saline climate, flooding.
Intensive pasture	Not suitable	Profuse gravels, saline climate, flooding.
Forestry	Not suitable	Profuse gravels, saline climate, flooding.

### **Management practices to improve suitability**

- None recommended.



**Whangaparoa Valley with Waiapu soils in the foreground, Oweka soils behind that, and Mataheiiia soils in the background flats on which houses and sheds have been built. Potikirua stepland soils are in the background**

## Soil Series Name: Waihoata (Wao)

### Overview

Waihoata soil series occur in small areas of low river flats south of Cape Runaway. Parent materials are alluvium derived from greywacke, mudstone and sandstone. Soil profiles consist of dark brown friable silt loam on greyish-brown clay loam with many yellowish-red mottles overlying light olive grey mottled clay loam. The soils are classified as **Acidic Recent Gley Soils**. Land use consists of grazing dry stock.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** 30 – 40 cm

**Rooting barrier:** Anoxic conditions

**Drainage class:** Poorly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Moderate to low (62 mm)

**Profile readily available water (0 – 100 cm):** Low (39 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.0 – 13.8%

**Topsoil P retention:** Medium (35%)

**Available P, Ca, Mg and K:** Possibly medium to low

### Soil types/variations

**Waihoata silt loam** as described above. Profiles with peaty layers and peaty loam topsoils occur.

#### ***Associated and similar soils***

**Oweka series** on slightly elevated terraces have well drained profiles. **Waiapu series** occur adjacent to stream beds with well drained and stony profiles.

## **General land use suitability ratings**

### **Waihoata silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Poor natural drainage, danger of flooding, small areas.
Horticulture	Not suitable	Poor natural drainage, danger of flooding, small areas.
Intensive pasture	Low	Poor natural drainage, danger of flooding.
Forestry	Not suitable	Poor natural drainage, danger of flooding.

### ***Management practices to improve suitability***

- Employ artificial drainage (open drains).



**Waihoata silt loam** (Note mottling in the subsoil)

## Soil Series Name: Waihurua (Ws)

### Overview

Waihurua soil series occur on dry watercourses south of the Rotorua–Waikaremoana Road, west of Murupara. Parent materials are alluvium derived from water-sorted pumice mostly derived from Taupo Tephra. Soil profiles consist of black gravelly coarse sand overlying dark brown and yellowish-brown loose gravelly sand on grey coarse gravelly sand. The soils are classified as **Immature Orthic Pumice Soils**. Land use consists of rough grazing and forestry.

### Physical properties

**Texture:** Skeletal

**Topsoil clay content:** 0%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (44 mm)

**Profile readily available water (0 – 100 cm):** Low (32 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Possibly low

### Soil types/variations

**Waihurua sand** has few lapilli in the topsoil. On the other hand, **Waihurua gravelly sand** has abundant coarse lapilli in the topsoil.

#### ***Associated and similar soils***

**Pekepeke series** are on adjacent rolling to hilly land.

## **General land use suitability ratings**

### **Waihurua sand and Waihurua gravelly sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Coarse textures with low water-holding capacity, cool climate.
Horticulture	Not suitable	Coarse textures with low water-holding capacity, cool climate.
Intensive pasture	Low	Coarse textures with low water-holding capacity, cool climate.
Forestry	Low	Coarse textures with low water-holding capacity.

### ***Management practices to improve suitability***

- Use zero or minimum tillage methods to preserve topsoil organic matter.
- Avoid cultivating the soil in dry, windy conditions.



**Waihurua sand**

## Soil Series Name: Waikaremoana (YkS)

### Overview

Waikaremoana soil series occur on steep and very steep slopes in and around Waikaremoana in Urewera National Park. Parent materials consist of rhyolitic tephra (very thin Kaharoa Tephra, and thin Taupo Tephra) on weathered rhyolitic tephra on hard sandstone or siltstone. Soil profiles are generally shallow and consist of dark brown fine sandy loam on olive yellow fine sandy loam on weathered sandstone. The soils are classified as **Typic Rocky Recent Soils**. Land use consists of hunting in indigenous forest.

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 10 – 15%

**Potential rooting depth:** 5 – 45 cm

**Rooting barrier:** Weathered fractured rock

**Drainage class:** Moderately well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Low (46 mm)

**Profile readily available water (0 – 100 cm):** Low (36 mm)

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 - 6.9 %

**Topsoil P retention:** Low (30%)

**Available P, Ca, Mg and K:** Low (strongly leached soils)

### Soil types/variations

**Waikaremoana steepland soils** are generally shallow soils on steep and very steep slopes. Kaharoa Tephra varies in thickness from 0 – 9 cm while Taupo Tephra varies from 10 – 30 cm. Total tephra thickness varies from 5 – 100 cm.

#### ***Associated and similar soils***

**Urewera steepland soils** are on steep and very steep slopes derived from tephra on greywacke, generally with thicker tephra. **Matawai hill soils** are on hilly slopes with thicker tephra.

## **General land use suitability ratings**

### **Waikaremoana steepland soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Not suitable	Steep and very steep slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low	Steep and very steep slopes, cool or cold climate.
Forestry	Low	Steep and very steep slopes, cold climate, erosion potential.

### **Management practices to improve suitability**

- These soils are best left in their natural state to avoid further erosion.
- Use aerial harvesting methods for forestry.
- Employ low stocking rates for dry stock.



**Waikaremoana steepland soil landscape**

## Soil Series Name: Waioeka (Woe)

### Overview

Waioeka soil series occur on former back swamps of the Waioeka river flood plain of the Opotiki flats. Parent material is alluvium derived from greywacke and rhyolitic tephra with 10 to 15 cm layers of diatomaceous earth and very thin layers of Kaharoa and Taupo Tephra. Soil profiles have deep dark brown friable silt loam on pale olive firm heavy silt loam with abundant mottles, resting on dark brown fine sandy loam on white hard silt (diatomaceous earth) which overlie pale yellow pumiceous sand (Taupo Pumice). Lower subsoils are brown mottled clay loam and greyish-brown clay. The soils are classified as **Acidic Orthic Gley Soils**. Land use is dairying and limited cash cropping (maize).

### Physical properties

**Texture:** Loam

**Topsoil clay content:** 20 – 25%

**Potential rooting depth:** 60 – 90 cm

**Rooting barrier:** Anoxic conditions

**Drainage class:** Poorly drained

**Permeability:** Slow

**Profile total available water (0 – 100 cm):** Moderate (72 mm)

**Profile readily available water (0 – 100 cm):** Moderate (60 mm)

**Topsoil bulk density:** 0.94 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.22 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 6.9 – 15.5%

**Topsoil P retention:** Medium (38%)

**Available P, Ca, Mg and K:** Low (Strongly leached soil)

### Soil types/variations

**Waioeka silt loam** as described above. Diatomaceous layers vary in thickness from 1 to 6 to 28 cm.

#### ***Associated and similar soils***

**Apanui series** are soils without the diatomaceous layers. **Otara series** are imperfectly drained and occur on the margins of the swamps.

## **General land use suitability ratings**

### **Waioeka silt loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Medium	Poor drainage, low fertility, heavy soil textures.
Horticulture	Low to medium	Poor drainage, low fertility, heavy soil textures.
Intensive pasture	Medium	Poor drainage, low fertility, pugging in winter.
Forestry	Not suitable	Poor drainage.

***Management practices to improve suitability***

- Employ artificial drainage (open or mole).



**Waioeka silt loam**

## Soil Series Name: Waipahihi (Yp)

### Overview

Waipahihi soil series occur in the pumice country on valley floors. Parent material is colluvium or alluvium derived from Taupo Tephra that eroded off adjacent hill slopes. Soil profiles have thin dark brown friable loamy sand topsoils with few rounded lapilli, on dark reddish-brown to yellowish-brown firm loamy sand with many rounded pumice lapilli. They overlie light yellowish-brown firm loamy sand and pale brown sand. The soils are classified as **Immature Orthic Pumice Soils**. Land use is dry stock and some dairying.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 5 – 8%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate to low (85 mm)

**Profile readily available water (0 – 100 cm):** Moderate (59 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.00 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%.

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Waipahihi loamy sand** as described above. **Waipahihi sand** has coarser textures towards the Taupo area.

#### ***Associated and similar soils***

**Taupo series, Oruanui series and Tauhara series** occur on adjacent rolling and hill country. **Mokai series** occur in low-lying swamps.

## **General land use suitability ratings**

### **Waipahihi loamy sand and Waipahihi sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Low	Coarse textures, cool climate, weakly developed topsoil structure, fodder cropping.
Horticulture	Low to unsuitable	Coarse textures, cool climate, weakly developed topsoil structure.
Intensive pasture	Medium to low	Coarse textures, cool climate, low fertility.
Forestry	Medium to low	Small areas.

### ***Management practices to improve suitability***

- Increase organic matter.
- Avoid soil tillage under dry, windy conditions.



**Waipahihi sand** (Note shallow topsoil and water-sorted layering in the subsoil, scale in m)

## Soil Series Name: Whakatane (Wx)

### Overview

Whakatane soil series occur on flat, rolling and hilly lands in the Whakatane area. Parent materials are less than 7 cm Tarawera Tephra on Kaharoa Tephra, Taupo Tephra, Whakatane Tephra and Rotoma Tephra. Soil profiles have very dark brown friable loamy sand on yellowish-brown friable loamy sand, resting on more yellowish-brown loamy sand. The soils are classified as **Allophanic Orthic Pumice Soils**. Land use is dairying, dry stock, cropping (maize), and forestry.

### Physical properties

**Texture:** Sand

**Topsoil clay content:** 3 – 10%

**Potential rooting depth:** Unlimited

**Rooting barrier:** No significant barrier within 1 m

**Drainage class:** Well drained

**Permeability:** Rapid

**Profile total available water (0 – 100 cm):** Moderate (96 mm)

**Profile readily available water (0 – 100 cm):** Moderate (72 mm)

**Topsoil bulk density:** 1.18 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.42 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%.

**Topsoil P retention:** Medium (51%)

**Available P, Ca, Mg and K:** Low

### Soil types/variations

**Whakatane loamy sand** are on easy rolling to rolling land. **Whakatane hill soils** are on hilly land. Thicker Tarawera Tephra and Kaharoa Tephra occur towards the western extent of the soils. Such profiles have thicker and blacker topsoils.

### *Associated and similar soils*

**Matahina series** are soils where Tarawera Tephra is more than 7 cm thick and coarser.

**Tawhia series** are on steep and very steep slopes with thinner, often mixed tephra, which overlies greywacke.

## **General land use suitability ratings**

### **Whakatane loamy sand**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Medium	Rolling slopes, weakly developed topsoil structure, low fertility.
Horticulture	Medium	Rolling slopes, weakly developed topsoil structure, low fertility.
Intensive pasture	Medium	Low fertility.
Forestry	High	No limitations.

### **Whakatane hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Unsuitable	Hilly slopes, weakly developed topsoil structure, low fertility.
Horticulture	Unsuitable	Hilly slopes, weakly developed topsoil structure, low fertility.
Intensive pasture	Medium	Hilly slopes.
Forestry	High	No limitations.

***Management practices to improve suitability***

- Avoid soil tillage under dry, windy conditions.



**Whakatane loamy sand**

## Soil Series Name: Whangaparoa (WhS)

### Overview

Whangaparoa soil series occur on steep and very steep slopes of a small area chiefly south of Cape Runaway and east of Whangaparoa Bay. Parent materials are coarse sandstone. Soil profiles have dark brown friable sandy loam on brownish-yellow sandy clay loam, resting at about 40 cm on yellow, coarse sandstone. The soils are classified as **Typic Orthic Recent Soils**. Land use is dry stock or forestry.

### Physical properties

**Texture:** Loam over sand

**Topsoil clay content:** 15 – 20%

**Potential rooting depth:** 40 – 60 cm

**Rooting barrier:** Weathered rock

**Drainage class:** Well drained

**Permeability:** Rapid over moderate

**Profile total available water (0 – 100 cm):** Moderate to low (74 mm)

**Profile readily available water (0 – 100 cm):** Low (47 mm))

**Topsoil bulk density:** 1.09 g/cm<sup>3</sup>

**Subsoil bulk density:** 1.30 g/cm<sup>3</sup>

### Chemical properties

**Topsoil organic matter:** 4.3 – 6.0%

**Topsoil P retention:** Low (22%)

**Available P, Ca, Mg and K:** Probably low

### Soil types/variations

**Whangaparoa steepland soils** are on steep and very steep slopes. Slightly deeper profiles occur on ridges and spurs.

#### *Associated and similar soils*

**Matakoa series and Wharekahika series** are soils on adjacent flat, rolling and hilly slopes.

#### *General land use suitability ratings*

##### Whangaparoa steepland soils

Land use	Suitability rating	Management considerations
Arable	Not suitable	Steep and very steep slopes.
Horticulture	Not suitable	Steep and very steep slopes.
Intensive pasture	Low to not suitable	Steep and very steep slopes, severe erosion potential.
Forestry	Low	Steep and very steep slopes.

### ***Management practices to improve suitability***

- Employ low stocking rates (sheep) to avoid accelerated erosion.
- Use aerial harvesting techniques for forestry.



**Whangaparoa steepland soil**

## Soil Series Name: Wharekahika (Wha)

### Overview

Wharekahika soil series occur on flat, rolling and hilly land in the Whangaparoa Catchment area, south of Cape Runaway. Parent materials are rhyolitic tephra on weathered basalt. Soil profiles have dark brown friable sandy loam on brown friable sandy loam, which rests at 48 cm depth on light yellowish-brown firm, mottled clay loam overlying olive yellow mottled clay. The clay is at about 16 cm depth on hilly slopes. The soils are classified as **Typic Orthic Brown Soils**. Land use is dairying, dry stock and limited cropping (maize).

### Physical properties

**Texture:** Loam over clay

**Topsoil clay content:** 10 – 15% (sandy loam); 25 – 30% (hill soils)

**Potential rooting depth:** Unlimited (sandy loam); 50 cm (hill soils)

**Rooting barrier:** No significant barrier within 1 m

**Stiff clay subsoils** Hill soils

**Drainage class:** Moderately well drained to imperfectly drained

**Permeability:** Moderate over slow

**Profile total available water (0 – 100 cm):** Moderate to high (153 mm) (hill 128 mm)

**Profile readily available water (0 – 100 cm):** Moderate (88 mm) (hill 70 mm)

**Topsoil bulk density:** 0.91 g/cm<sup>3</sup> (hill 1.09 g/cm<sup>3</sup>)

**Subsoil bulk density:** 0.84 g/cm<sup>3</sup> (hill 1.26 g/cm<sup>3</sup>)

### Chemical properties

**Topsoil organic matter:** 8.6 – 12.1%.

**Topsoil P retention:** Medium (36 – 51%)

**Available P, Ca, Mg, S and K:** Low to medium

### Soil types/variations

**Wharekahika sandy loam** occurs on flat to rolling surfaces. **Wharekahika hill soils** occur on hilly slopes.

#### *Associated and similar soils*

**Matakaoa series** are well drained soils with similar parent materials.

## **General land use suitability ratings**

### **Wharekahika sandy loam**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Medium	Rolling slopes, imperfectly drained areas.
Horticulture	Low	Rolling slopes, imperfectly drained areas.
Intensive pasture	Medium	Imperfect drainage, pugging risk, strong erosion potential.
Forestry	Low	Imperfect drainage, strong erosion potential.

### **Wharekahika hill soils**

<b>Land use</b>	<b>Suitability rating</b>	<b>Management considerations</b>
Arable	Unsuitable	Hilly slopes, imperfectly drained areas.
Horticulture	Unsuitable	Hilly slopes, imperfectly drained areas.
Intensive pasture	Medium	Hilly slopes, imperfect drainage: pugging risk, strong erosion potential.
Forestry	Low	Hilly slopes, imperfect drainage, strong erosion potential.

### ***Management Practices to Improve Suitability***

- Avoid soil tillage under dry, windy conditions.
- Avoid high stocking rates because of erosion risk especially on hilly slopes.
- Avoid high stocking rates because of high pugging risk under wet conditions.



**Wharekahika sandy loam**



## Part 8: Glossary

---

**ABC soil** A soil that has an A, a B, and a C horizon.

**Absorption** Filling up of soil pores with water like a sponge soaks up water.

**AC soil** A soil that only has an A and a C horizon. Commonly, such soil formed in recent alluvium or on steep, rocky slopes.

**Adsorption** The attraction of ions or compounds to the surface of a solid. Soil colloids (clay and humus) adsorb large amounts of ions and water.

**Aeolian** Refers to wind-blown soil materials such as loess and sand.

**Aeration** The movement of air back and forth between the atmosphere and the pores of a soil.

**Allophane** A non-crystalline hydrous aluminium silicate clay mineral found in volcanic ash soils.

**Allophanic** Soils that are dominated by the clay mineral allophane (and also imogolite, ferrihydrite, and/or aluminium-humus complexes). They have a characteristically greasy feel, and high to very high phosphate retention.

**Alluvium** General term for unconsolidated materials such as gravel, sand, silt, clay or mixtures of these deposited on land by streams.

**Andesite** Volcanic rock composed essentially of andesine and one or more mafic constituents (such as pyroxene, hornblende, or biotite).

**Anoxic** Lacking oxygen, as in waterlogged soils.

**Available water capacity (or available water-holding capacity)** The maximum amount of water a soil can store for plant use in the upper 1 m. It is the difference between the amount of soil water held at **field capacity** (-10 kPa) and the amount at **wilting point** (-1,500 kPa). It is commonly expressed as mm of water. A good soil can provide around 230 – 305 mm of available water, a poor soil about 50 – 75 mm only.

**Base saturation** The extent to which the adsorption complex of a soil is saturated with exchangeable cations other than hydrogen and aluminium. It is expressed as a percentage of the soil's **cation exchange capacity**.

**Breccia** A coarse grained, clastic rock composed of angular, broken rock fragments held together by a mineral cement or a fine-grained matrix.

**Boulders** Rock fragments larger than 200 mm in diameter.

**Buff layer** Compact layer of alluvium derived from Kaharoa Tephra present in some soils such as Awaiti, Omeheu and Paroa series.

**Bulk density** Mass of dry soil per unit of bulk volume where **bulk volume** includes the volume of solids as well as the volume of pore spaces (total volume). It is expressed in g/cm<sup>3</sup> or t/m<sup>3</sup>.

**Caldera** A large basin-shaped volcanic depression, more or less circular in form.

**Cation exchange capacity** The sum total of exchangeable cations that a soil can adsorb. It is expressed as centimoles of charge per kilogram of soil (cmol/kg).

**Clastic** Consisting of fragments of rocks.

**Clay** (1) Mineral soil particle with a diameter of less than 0.002 mm; (2) A soil textural class which is very plastic, very sticky and contains 60% or more clay.

**Colour** Colour gives some clues about the soil. Dark topsoils usually indicate high organic matter levels. Grey colours are indicative of poorly-drained soils. Yellowish-brown and reddish-brown colours generally indicate favourable air – water relations.

**Colluvium** Rock fragments or soil materials that have accumulated at the base of steep slopes due to gravity.

**Compaction** Reduction in the porosity of a soil due to cultivation and other mechanical forces.

**Consistence** The degree of cohesion and adhesion of peds, and the ease with which they are dislodged from the soil profile.

**Creep (also called scree creep)** Slow mass movement of soil down relatively steep slopes, mainly under the influence of gravity, but enhanced by saturation with water and by alternate freezing and thawing. Commonly seen as terracettes on hill slopes.

**Dacite** Extrusive rock with principal minerals plagioclase (andesine and oligoclase), quartz, pyroxene or hornblende or both, with minor biotite and sanidine.

**Diatomaceous earth** Fine, greyish siliceous material composed primarily of the remains of diatoms that have accumulated mainly in water. It may occur as a powder or as a porous, rigid material.

**Drainage class** The degree of wetness of a soil, as determined by the depth to a water table and the length of time the soil remains saturated. Common drainage classes include well-drained, moderately well-drained, imperfectly drained and poorly-drained.

**Droughty** A soil incapable of storing much water for plant use. Coarse sandy, stony, gravelly, sandy, shallow and steeply sloping soils are likely to be droughty unless they are supplied with rainfall or irrigation water frequently.

**Dune** A low hill or bank of drifted sand. Also mounds or ridges of wind-blown or aeolian sand are dunes.

**Evapotranspiration.** Between periods of rain, water held in the soil is gradually given up by direct evaporation from the soil surface and by plant transpiration. The combined water loss by these two means is called evapotranspiration.

**Fan (alluvial)** A low, outspread mass of loose materials and/or rock material, commonly with gentle slopes, shaped like an open fan or a segment of a cone, deposited by a stream.

**Field capacity** The moisture content of a soil when free drainage (one to two days after a heavy rain or irrigation) has virtually stopped. It is the maximum amount of water a soil can retain against the force of gravity. It is also called the upper limit of available water.

**Fine earth** All soil materials less than 2 mm in diameter.

**Flats** A flat or surface of low relief. Generally used to indicate flat river terraces.

**Floodplain** The nearly level surface next to a river that is covered with water when the river floods.

**Gleyed** Soil that is very wet for long periods of time and is characterized by grey colours, with or without mottles.

**Gley soil** Soil that formed under poor drainage, resulting in the reduction of iron, manganese and other elements in the profile and in grey colours.

**Gravel** Small, rounded coarse fragments in soils with a diameter ranging from 2 mm – 200 mm.

**Greywacke** A type of sandstone characterised by its dark colour, hardness and angular rock particles in a clayey matrix.

**Groundwater** The water below ground which saturates the subsoil. The upper surface of the zone of saturation is the water table.

**Hummocky** Regular or irregular small elevations or hillocks.

**Humus** The relatively resistant fraction of soil organic matter that forms during biological decomposition of organic residues. Humus usually constitutes the major fraction of soil organic matter.

**Ignimbrite** An igneous rock formed by the lithification of ash flow or pyroclastic flow deposits.

**Inclusion** An area of soil that is too small to show separately at the scale of the soil map. Inclusions can only be mapped out separately by making very detailed maps at very large scales.

**Infiltration** The rate at which water enters the soil. It is dependent on the size of pores and the stability of soil aggregates on the soil surface. If water cannot infiltrate, it either ponds on the surface or runs off over the surface.

**Landform** Any recognisable physical form or feature of the earth's surface having a characteristic shape and resulting from natural causes.

**Landscape** A portion of the land that the eye can comprehend in a single view, including all its natural characteristics.

**Lapilli** See **tephra**.

**Leaching** Removal of soluble materials such as minerals, nutrients, organic chemicals and pesticides from the soil by water passing through it either as rainfall or irrigation.

**Levee** A long, broad low ridge or embankment of sand and coarse silt, built by a stream on its flood plain and along both sides of its channel, especially in time of flood when water overflowing the normal banks is forced to deposit the coarsest part of its load.

**Limiting layer** Gravel, pan or stagnant water that limits plant root growth.

**Loess** Silt and fine sand particles deposited by the wind.

**Mottles or Mottling** Spots or blotches of grey or brown colour different from the dominant soil colour that indicate the height of fluctuating water tables and hence the degree of soil aeration.

**Muck** See **peat**.

**Mudstone** Silt- and clay-containing sedimentary rock that is non-plastic and has a massive appearance.

**Organic matter** Plant and animal residues in the soil in various stages of decomposition.

**Organic soil** Soil formed in the partly decomposed remains of wetland plants (peat) or forest litter. Some mineral material may be present but the soil is dominated by organic matter.

**Oxbow** A crescent-shaped lake formed in an abandoned river bend which has become separated from the stream by a change in the course of the river

**Paleosol** A soil that formed on a landscape in the past and that has distinctive morphological characteristics resulting from a soil-forming environment that no longer exists at the site. The former soil-forming process was either altered due to environmental change (e.g. climatic change) or interrupted by burial (e.g. occurrence of a **buried soil** during a volcanic eruption or past flooding).

**Pan** A compact, dense layer in a soil that restricts the movement of water and penetration by plant roots.

**Parent material** Unconsolidated organic and mineral material from which the soil develops.

**Parent rock** The rock from which the parent material is derived.

**Peat** Organic soil material in which the plant residues are still recognisable. This contrasts with **muck** in which the original plant residues cannot be recognised any more.

**Ped** A naturally-occurring soil aggregate, as opposed to a **clod**, which is formed artificially (e.g. through cultivation).

**Permeability** The rate at which water moves through the soil. It depends on the amount, size and interconnectedness of the pores, which in turn depend on soil texture, structure and bulk density.

**pH** The degree of acidity or alkalinity of a soil. A pH of 7 indicates a neutral soil. Lower values indicate acidic soils; higher numbers indicate alkaline soils.

**Phosphate, or P, retention** Expressed as a percentage, this is a measure of the degree of phosphate retention or immobilisation by soil minerals. For the same amount of phosphate fertiliser applied, soils with high P-retention values will give lower crop yields than soils with low P-retention values.

**Podzol** A strongly acidic soil that usually has a bleached soil horizon immediately below the topsoil. It occurs in high rainfall areas and is associated with forest trees with an acidic litter.

**Pore space** Soil space not occupied by solid particles (i.e. occupied by air and water).

**Porosity** The total volume of pore space in the soil expressed as a percentage of the bulk or total soil volume.

**Potential rooting depth** Total depth of soil suitable for plant root growth, measured from the surface to the top of a barrier (within 1 m of the soil surface) that limits root extension. Actual rooting depth depends on the depth plant roots actually penetrate.

**Pumice** Light vesicular form of volcanic glass with a high silica content. It is usually light in colour and some can float on water.

**Pumice soil** A soil with properties dominated by pumice and glass with a low clay content (which contains allophane). It occurs in sandy or pumiceous tephra with an age range of 700 – 3,500 years.

**Readily available water** The amount of water easily extractable by plant roots. It is the amount of water held between -10 kPa and -100 kPa suction. It is expressed as an equivalent depth of water in mm.

**Ridge** A relatively narrow elevation which is prominent on account of the steep angle at which it rises.

**Rooting barrier** The type of barrier that limits root penetration (e.g. compact soil horizons, pans, rocks, densely packed gravels, anaerobic conditions and high water tables).

**Rhyolite** A light, fined-grained igneous rock formed by the rapid cooling of lava rich in silica.

**Runoff** Water that flows over the surface of soil toward a stream or lake without sinking into the soil.

**Sand** (1) Mineral soil particle with a diameter range of 2.0 – 0.06 mm. Further subdivided into coarse sand (2.0 – 0.6 mm), medium sand (0.6 – 0.2 mm) and fine sand (0.2 – 0.06 mm); (2) A soil textural class containing 80% or more sand.

**Sandstone** Sedimentary rock consisting predominantly of sand-sized particles.

**Scree** A heap of rock waste at the base of a cliff or a sheet of coarse debris mantling a mountain slope.

**Seepage** Water that seeps toward stream channels after infiltration into the ground.

**Shale** Sedimentary rock formed by the hardening of a clay deposit.

**Silt** (1) Mineral soil particle with a diameter range of 0.06 – 0.002 mm; (2) A soil textural class (82% or more silt) which feels extremely smooth and silky.

**Siltstone** Sedimentary rock made up of predominantly silt-sized particles.

**Skeletal** Soils with horizons containing 35% or more gravel by volume.

**Slope** The inclination of the land surface expressed in degree angle from the horizontal.

**Soil** A natural body on the earth's surface that develops in response to climate and organisms acting on a parent material in a specific landscape position over a long period of time. Soil supplies plants with air, water, nutrients and provides mechanical support.

**Soil association** A group of soils geographically associated in a characteristic repeating pattern, and defined and delineated as a single map unit.

**Soil complex** (1) A map unit of two or more kinds of soil so intimately intermixed geographically that it is not practical to map them separately at the selected scale of mapping; (2) A more intimate mixing of smaller areas of individual soil mapping units than that in a soil association.

**Soil horizon** A layer of soil, approximately parallel to the surface, having distinct characteristics produced by soil-forming processes. From top to bottom, soil horizons are named as A, B, and C.

**Soil morphology** The physical makeup of the soil, which includes thickness and arrangement of soil horizons, colour, texture, structure, consistence, roots, horizon boundaries, etc.

**Soil phase** A subdivision of a soil series based on features that affect its use and management such as slope, stoniness, flooding, etc.

**Soil profile** A vertical section of the soil showing all its horizons and extending into the parent material.

**Soil series** A group of soils that have similar profile characteristics except for differences in texture of the surface layer. Traditionally, a soil series is named after the place where the soil was first observed and described (e.g. Taupo soil series).

**Soil survey** The process of producing a soil map. A soil surveyor walks over the land, and observes and records soil and landscape features, classifies the soils, and locates soil boundaries in the field. The surveyor uses aerial photos as base maps to delineate soil boundaries and label each area, or polygon, with a map unit symbol.

**Soil survey report** A publication containing the results of a soil survey of an area. It consists of the soil map, text describing the properties and behaviour of soils, and tables containing interpretations for soil use and management.

**Soil type** A subdivision within a soil series to distinguish soils differing in surface texture only (e.g. Taupo sand, Taupo sandy loam, etc.).

**Soil variant** A soil whose properties are believed to be different enough from other known soils to justify creation of a new series name but, because of its limited geographic extent, the creation is not justified.

**Stones** Rock fragments ranging from boulders to gravels.

**Stony** Soil that contains 5 – 35% stones in the upper 20 cm. Stones limit the volume of soil that is available for roots to explore for water and nutrients. They can also significantly interfere with or prevent tillage operations. Where stones consist of lapilli, these do not adversely affect land use since these are light materials and do not present an impediment to cultivation.

**Spur** A subordinate ridge which extends itself from the crest of a hill or mountain like ribs from the vertebral column.

**Structure** The arrangement of the primary particles sand, silt and clay into larger units called aggregates or peds. Plant roots, clay and organic matter help bind aggregates together.

**Subsidence** Lowering of the soil surface due to settling or shrinkage. Drainage of organic soils results in subsidence through increased aeration, and the loss of organic matter through decomposition.

**Swale** A slight depression in an area of generally flat land.

**Tephra** A name for all unconsolidated **clastic** volcanic material that, during an eruption, is transported through the air from the source. This term should not be confused with grain size classes: fine ash (less than 0.25 mm); coarse ash (0.25 – 2.0 mm); lapilli (2.0 – 64 mm).

**Texture** The relative proportion of sand, silt and clay particles in a soil. Specific combinations of sand, silt and clay are known as **textural classes**. Examples include sandy loam, silt loam, clay, etc.

**Vesicular** Containing many small cavities or air holes (as in Taupo Pumice).

**Volcanic ash** Fine, ash-like rock particles ejected from a volcano during an eruption that may be transported long distances by the wind.

**Water table** The upper level of water stored under the ground.

**Wilting point** The moisture content of a soil at which plants can no longer extract water. Clayey soils contain relatively large amounts of water at the wilting point, but it is held so tightly inside the very small pores of the clay that plants are unable to extract it.

**Years before present** Carbon-14 dating system for tephra using 1950 as the base year.



## Part 9: References

---

- Auckland Regional Council. 1999. Erosion and sediment control: guidelines for land disturbing activities in the Auckland Region. Auckland Regional Council Technical Publication No. 90.
- Blakemore, L.C., Searle, P.L. and Daly, B.K. 1987. Methods for Chemical Analyses of Soils. NZ Soil Bureau Scientific Report 80.
- Carrick, S. 2007. Preferential flow through soils. *Soil Horizons*, Issue 16 (September), p. 8. Available online at: <http://www.landcareresearch.co.nz/publications/newsletters/soilhorizons/SoilHorizIssue16Sept07.pdf>
- Cornforth, I. 1998. Practical Soil Management. Lincoln University Press and Daphne Brassel Associates, Ltd. Canterbury and Wellington, New Zealand. 248 p.
- Cotching, W. E. 1998. Soil Survey of the Te Puke District, Bay of Plenty, New Zealand. Landcare Research Technical Record. Manaaki Whenua – Landcare Research New Zealand Ltd, Hamilton, New Zealand.
- Froggatt, P.C. and Lowe, D.J. 1990. A review of late Quaternary silicic and some other tephra formations from New Zealand: their stratigraphy, nomenclature, distribution, volume and age. *New Zealand Journal of Geology and Geophysics* 33: 89-109.
- Griffiths, E. 2004. Soils of the Ruataniwha Plains: A Guide to Their Management. Grifftech and Hawke's Bay Regional Council, Napier.
- Hewitt, A. E. 1998a. New Zealand Soil Classification. 2<sup>nd</sup> edition. Landcare Research Science Series No. 1, Manaaki Whenua Press, Lincoln, Canterbury, New Zealand.
- Hewitt, A. E. 1998b. Appendix: The New Zealand soil classification, *In* D. Molloy. Soils in the New Zealand Landscape: The Living Mantle. 2<sup>nd</sup> ed. Mallinson Rendell in association with New Zealand Society of Soil Science, Wellington. 253 p.
- Hewitt, A.E. 2004. Soil properties for plant growth: a guide to recognizing soil attributes relevant to plant growth and plant selection. Landcare Research Science Series No. 26. Lincoln, Canterbury, New Zealand.
- Hicks, D. H. and Anthony, T. 2001. Soil Conservation Technical Handbook, Parts A and B. Ministry for the Environment, Wellington, New Zealand. Available online at: <http://www.mfe.govt.nz/publications/land/soil-conservation-handbookjun01/soil-conserv-handbook-jun01.pdf>
- Lowe, D. J. (ed.) 2006. Guidebook for 'Land and Lakes' field trip, NZ Society of Soil Science Biennial Conference, Rotorua. New Zealand Society of Soil Science, Lincoln, 63 pp.
- Lowe, D.J. and Palmer, D.J. 2005. Andisols of New Zealand and Australia. *Journal of Integrated Field Science* 2:39-65.
- Lynn, I. H., Manderson, A. K., Page, M. J. Harmsworth, G. R., Eyles, G. O., Douglas, G. B., Mackay, A. D. and Newsome, P. J. F. 2009. Land Use Capability Survey Handbook: A New Zealand Handbook for the Classification of Land. 3rd ed. Hamilton, Agresearch; Lincoln, Landcare Research; Lower Hutt, GNS Science. 163 p.

- Manderson, A., Palmer, A., MacKay, A., Wilde, H., and Rijkse, W. 2007. Introductory Guide to Farm Soil Mapping. AgResearch Ltd, Palmerston North, New Zealand.
- Milne, J.D.G., Clayden, B., Singleton, P. L. and Wilson, A. D. 1995. Soil Description Handbook. Revised edition. Manaaki Whenua Press, Lincoln, Canterbury, New Zealand.
- Molloy, D. 1998. Soils in the New Zealand Landscape: The Living Mantle. 2<sup>nd</sup> ed. New Zealand Society of Soil Science, 253 p.
- Morgan, R.P.C. 1995. Soil Erosion and Conservation. 2<sup>nd</sup> ed. Longman Group Limited, Essex, England.
- New Zealand Soil Bureau. 1954. General Survey of the Soils of the North Island, New Zealand. New Zealand Soil Bureau Bulletin 5.
- Ngapo, N. 2010. Erosion types in the Bay of Plenty / Volcanic Plateau, Eastern and Southern Waikato (Powerpoint notes). In "Soil Erosion Workshop" sponsored by Ministry of Agriculture and Forestry and Environment Bay of Plenty. 23 April 2010, SCION, Rotorua.
- Pullar, W. A. 1972. Soil Maps and Extended Legend of Whakatane Borough and Environs, Bay Of Plenty, New Zealand. New Zealand. Department of Scientific and Industrial Research. Soil Bureau.
- Pullar.W.A., Birrell, K.S. 1973 Age and Distribution of Late Quaternary Pyroclastic and Associated Cover Deposits of the Rotorua and Taupo Area, North Island, New Zealand. N.Z. Soil Survey Report 1 Parts 1, 2 and 3.
- Pullar, W. A. 1985. Soils and Land Use of Rangitaiki Plains, North Island, New Zealand. New Zealand Soil Bureau.
- Pullar, W.A., Hewitt, S.R. and Heine, J.C. 1978. Soils and Land Use of Whakatane Borough and Environs, Bay of Plenty, New Zealand. New Zealand Department of Scientific and Industrial Research.
- Rijkse, W. C. 1979. Soils of Rotorua Lakes District, North Island, New Zealand. NZ Soil Survey Report 43, New Zealand Soil Bureau, Department of Scientific and Industrial Research, Wellington, New Zealand.
- Rijkse, W. C. 1993. Soils of Whakatane County, North Island, New Zealand. Landcare Technical Research Record.
- Rijkse, W.C. 1997. Soils, Agriculture and Forestry of Taupo Region, North Island, New Zealand. Landcare Research New Zealand Scientific Report. Landcare Research New Zealand Ltd, Hamilton, New Zealand.
- Rijkse, W.C. 1988. Soils of the Kaingaroa Plateau, North Island, New Zealand. New Zealand Soil Bureau., Rotorua, New Zealand. 127 pp.
- Rijkse, W.C. 1997. Soils and Agriculture of Opotiki District, Bay of Plenty, New Zealand. Landcare Research Ltd, Hamilton, NZ 178 pp.
- Rijkse, W.C. 2001. Soils of the Whakarewarewa and Rotoehu Forests, Bay of Plenty, New Zealand. Landcare Research Ltd, Hamilton. 65 pp.
- Rijkse, W.C. 2001. Landcare Research Contract Report 0001/148.Landcare Research New Zealand Ltd, Hamilton, New Zealand.

- Rijkse, W.C. 2002. Soils and Land Use of Katikati Area, Bay of Plenty, New Zealand. Landcare Research Ltd, Hamilton, NZ. 70 pp.
- Rijkse, W. C. 2002. Soils and Land Use of Part Topographic Sheets U15 and V15, Bay of Plenty, New Zealand. Landcare Research Technical Record. Landcare Research New Zealand Ltd, Hamilton, New Zealand.
- Rijkse, W.C. 2002. Soils of Mamaku and Ngamanawa Forests, Bay Of Plenty, New Zealand. Landcare Research Contract Report LC 0203/032. Landcare Research New Zealand Ltd, Hamilton, New Zealand.
- Rijkse, W.C. and Cotching, W.E. 1995. Soils and land use of part Tauranga County, North Island, New Zealand. Landcare Research Technical Record.
- Rijkse, W.C. 2004. Soils and Land Use of Eastern Bay of Plenty, New Zealand. Land Use Consultants Report BOP 1. Land Use Consultants, Mount Maunganui.
- Rijkse, W.C. 2004. Soils of Paengaroa Forests, Bay of Plenty, New Zealand. Landcare Research Contract Report LC 0304/076. Landcare Research New Zealand Ltd, Hamilton, New Zealand.
- Rijkse, W. C. 2004. Soils of Tarawera and Rotoiti Forests, Bay of Plenty, New Zealand. Land Use Consultants Ltd, Mount Maunganui.
- Shepherd, T. G. 2009. Visual Soil Assessment, Volume 1: Field guide for pastoral grazing and cropping on flat to rolling country. 2<sup>nd</sup> ed. Horizons Regional Council, Palmerston North. 119 p.
- Trought, M. C. T. 1981. Root aeration and the accumulation of nutrients by plants. *New Zealand Agricultural Science* 15:182.
- University of Minnesota 2000. Soil Management Series, Part 1: Soil Management. University of Minnesota Extension Publication BU-7399. ([http://www.extension.umn.edu/distribution/cropsystems/components/7399\\_02.html](http://www.extension.umn.edu/distribution/cropsystems/components/7399_02.html))
- U.S. Department of Agriculture, Natural Resources Conservation Service. National Soil Survey Handbook. Glossary of landform and geologic terms Part 629, pages 629-1 to 629-127. Available online at: <http://soils.usda.gov/technical/handbook>.
- Vucetich, C.G., Leamy, M.L., Popplewell, M.A., Ure, J., Taylor, C.R., Will, G.M., Sutton, J. A. and Blakemore, L. C. 1960. Soils, Forestry and Agriculture of the Northern Part, Kaingaroa State Forest and the Galatea Basin. Soil Bureau Bulletin 18. New Zealand Department of Scientific and Industrial Research.
- Vucetich, C.G. and Wells, N. 1978. Soils, Agriculture and Forestry of Waiotapu Region, Central North Island, New Zealand. NZ Soil Bureau Bulletin 31.



## Part 10: Index to Eastern Bay of Plenty soil series

No.	Soil Series Name	Map Symbol	Soil Order	Page
1	Amokura	Am	Gley	37
2	Apanui	Ap	Gley	39
3	Awaiti	Ati	Recent	41
4	Awakaponga	Ag	Recent	43
5	Awakeri	Aw	Pumice	46
6	Galatea	G	Pumice	49
7	Hanaia	Hi	Gley	52
8	Horomanga	Hm	Recent	55
9	Iwiroa	Ir	Gley	58
10	Kawerau	Kr	Pumice	60
11	Kopeopeo	Koe	Podzol	63
12	Kopuriki	Kpu	Pumice	66
13	Kukumoa	Ku	Recent	68
14	Manawahe	Mj	Pumice	71
15	Mangaomeko	MaH	Brown	74
16	Man-made	MM	Anthropic	77
17	Mataheia	Map	Recent	79
18	Matahina	Mb	Pumice	82
19	Matakaoa	Mak	Allophanic	85
20	Matao	Ma	Organic	88
21	Matata	Maa	Recent	90
22	Matawai	Mx	Podzol	93
23	Matuku	Mtk	Gley	96
24	Motu	MotH	Brown	99
25	Ngatiawa	NS	Recent	102
26	Ohope	Oe	Raw	104
27	Okoia	Ok	Gley	107
28	Omarumutu	Om	Gley	109
29	Omeheu	Ome	Gley	111
30	Onepu	Onp	Gley	114
31	Opotiki	Op	Pumice	117
32	Opouriao	Ou	Recent	120
33	Opoutama	Ox	Raw	122
34	Orangihewa	Og	Pumice	125

No.	Soil Series Name	Map Symbol	Soil Order	Page
35	Oruanui	Oi	Pumice	128
36	Otara	Ota	Recent	131
37	Oweka	Ow	Recent	133
38	Paerata	Pe	Gley	135
39	Paroa	Pr	Gley	137
40	Piripai	Pi	Raw	139
41	Pongakawa	Po	Organic	141
42	Poronui	Poi	Pumice	143
43	Potikirua	PotS	Recent	145
44	Pukemaku	PkS	Pumice	147
45	Pukerimu	Py	Pumice	149
46	Pukeroa	Pu	Pumice	152
47	Pureora	Pur	Podzol	154
48	Rangitaiki	Ran	Recent	156
49	Raukumara	RauS	Recent	158
50	Rewatu	Re	Gley	161
51	Ruakituri	Ru	Pumice	164
52	Ruatoki	Rti	Recent	167
53	Taho	Ta	Gley	170
54	Takamore	Tsl	Recent	172
55	Tarawera	Tr	Recent	174
56	Tauhara	TaS	Pumice	177
57	Taupo	Tp	Pumice	179
58	Tawhia	TyS	Pumice	182
59	Te Kaha	TeK	Brown	184
60	Te Piki	Pik	Organic	186
61	Te Rahu	TRa	Pumice	189
62	Te Teki	TtS	Podzol	191
63	Te Teko	Tto	Pumice	193
64	Tihoi	Toi	Podzol	195
65	Tokata	To	Gley	198
66	Torere	Tor	Raw/Recent	200
67	Tuparoa	TupS	Recent	202
68	Tutaetoko	TutS	Recent	204
69	Urewera	US	Pumice/Recent	206
70	Waiapu	Wu	Recent/Raw	208
71	Waihoata	Wao	Gley	210

<b>No.</b>	<b>Soil Series Name</b>	<b>Map Symbol</b>	<b>Soil Order</b>	<b>Page</b>
72	Waihuru	Ws	Pumice	213
73	Waikaremoana	YkS	Recent	216
74	Waioeka	Woe	Gley	218
75	Waipahihi	Yp	Pumice	221
76	Whakatane	Wx	Pumice	224
77	Whangaparoa	WhS	Recent	227
78	Wharekahika	Wha	Brown	229