



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Manawahe Kokako Survey Report

November 2014



Photo by Bruce Bancroft

Prepared for Bay of Plenty Regional Council and the Manawahe Kokako Trust

By Bruce Bancroft and Nicole Bancroft



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Table of Contents

Introduction.....	3
Methods.....	3
Results.....	4
Table 1. Kokako Population Comparisons 2013/2014	4
Other Birds.....	4
Pest Control.....	5
Discussion	5
Table 2. Manawahe Kokako Survey Results Over Time.....	7
Table 3. Manawahe Kokako Pairs change over time.....	8
Recommendations.....	9
References:.....	9
Graph 1. Manawahe Kokako Pair Total/Pair Trend Comparison 2014.....	12
Graph 2. Manawahe Kokako Pairs, Rat Tracking Data & Possum RTC Over Time	13
* Kokako pair only – double for actual number.	13
Map 1. Manawahe Kokako Area Surveyed 2013	14
Map 2. Kokako Pair Territory Comparison 2012-2013.....	15
Map 3. Kokako habitat around survey area	16
.....	16



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Manawahe Kokako Survey Report

November 2014

Introduction

The Manawahe Kokako Trust area and surrounds were resurveyed between September and November 2014 to establish the current status of the Manawahe kokako population. The previous census was conducted between October and November 2013.

Methods

Kokako are territorial birds, therefore calls and song are important if kokako are to retain their territories. Territory mapping is an essential tool in ascertaining population size.

Pre-recorded local kokako dialect was used during the survey to attract territorial kokako and was played through a portable speaker via MP3 digital player.

Two observers were used in tandem during the survey. To accurately determine kokako territorial boundaries, one observer would stay with observed kokako and follow them while the other would walk ahead playing kokako audio. Invariably kokako would follow the audio to the edge of their territory where they would remain and this is often when neighbouring kokako would turn up nearby, hence the need for two observers.

Both observers would communicate with hand held radios to compare sightings, this ensured that no double counting or under counting occurred.

GPS tracking data and the proximity of kokako to bait station sites were used to construct territory maps on GIS mapping software.



Eco Rescue Ltd
Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Results

The 2013 survey established the presence of 15 pair and 7 single kokako, giving a total population of 37 kokako.

The 2014 survey established the presence of 13 pair and 7 single kokako, a total population of 33 kokako.

The single present in the Cell-phone tower block to the north is still single.

Table 1. Kokako Population Comparisons 2013/2014

2012/2013 Kokako Population Comparisons				
	Pairs	Singles	Juveniles	Totals
2013	15	7	0	37
2014	13	7	0	33
Difference	-2	0	0	-4

Other Birds

Tui and bellbird are still in very good numbers as are kereru. Robins and pied tit seem to have increased considerably.

Still of concern are the large number of magpies seen and heard throughout all of the area surveyed. Harrier hawks were also noticeable throughout the survey. Myna are now present after not being noticed in previous surveys.



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Pest Control

During 2013, prior to the bait station toxin operation, gaps identified in the overall bait station layout were filled in with additional bait stations. This along with the predator control implemented from 2009 means the 2013-2014 breeding season would have the most effective pest control coverage in the operational area since the start of this project.

The Manawahe Kokako Trust carried out their annual bait station filling operation with 500g Pindone and 2 Feratox on 7th September 2013. This achieved a 0% rat tracking index on the five 10 tunnel tracking tunnel lines in the main kokako block as measured on 15th November 2013. This is 0% at 9 weeks after bait was laid and is a typical result that we regularly measure at 6 weeks at other sites. It is also the first time since 2008 this target has been met prior to the kokako breeding season.

Discussion

The overall population has fallen by four birds since the last survey; the number of pairs is down two and singles remain the same. Some territories were surveyed two or three times on separate days to minimise the chance of over or undercounting. This included some single bird territories to ascertain whether they were in fact single or a pair (female may have been sitting on an early nest). Care was taken to ensure all kokako in the survey area were accounted for in this survey but there always remains the possibility of birds, especially lone birds, not responding to the audio.

Degradation of the understory was noticeable in some areas within the survey boundary, probably due to deer and wallaby browsing. This may also be impacting on food availability. Pig sign was minimal.



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

The following paragraph is taken from the Manawahe Kokako Survey Report 2010. It has been repeated in each report since to remind readers of the main issues associated with managing a small remnant kokako population.

“Young birds may travel substantial (considering their limited flight ability) distances but generally settle near other kokako (Innes et al, unpubl. data). This behaviour is not surprising. Research on several species suggests the presence of conspecifics often attracts dispersing juveniles; the presence of adults may indicate that the area contains potential mates, food, nesting sites, or other important resources (Ahlering & Faaborg 2006). Because kokako are much more conspicuous vocally than they are visually, it is likely that their patterns of dispersal are influenced by the song of conspecifics.” (Calcott et al., 2008)

It has been suggested that birds may be dispersing to areas away from the main operational area. In previous surveys some of these outlying areas were surveyed with no kokako encountered. My experience with other kokako populations in Te Urewera suggests that kokako do not travel far from other kokako.

Also to consider: - *“In essence, at very low population levels, widely dispersed individuals have difficulty in finding mates and hence their reproduction should be lower than those living in higher numbers (Dennis, 1989; Courchamp et al., 1999; Stephens and Sutherland, 1999).”*

Table 1 and Graphs 1 and 2 show the population growth and decline in graphic form over the period of active management from the first census in 1997 through to the latest pre breeding census in October 2014.



Eco Rescue Ltd
 Bruce Bancroft
 0276318599
 bruce@ecorescue.co.nz

Table 2. Manawahe Kokako Survey Results Over Time

Manawahe Kokako Survey Results				
	Pairs	Singles	Juveniles	Totals
May 1997				14
May 2001	8	4	4	24
Apr 2002	10	2	10	32
May 2003	14	3	3	34
May 2004	19	2	4	44
May 2005	21	2	7	51
Oct 2007	24	5	0	53
May 2009	16	6	4	42
Oct 2010	17	4	0	38
Oct 2011	17	6	0	40
Oct 2012	16	8	0	40
Oct 2013	15	7	0	37
Oct 2014	13	7	0	33



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Table 3. Manawahe Kokako Pairs change over time.

	Pairs	Difference	% Change
May-2001	8		
Apr-2002	10	2	25%
May-2003	14	4	40%
May-2004	19	5	36%
May-2005	21	2	11%
Jun-2006	23	2	10%
Oct-2007	24	1	4%
Jun-2008	20	-4	-17%
May-2009	16	-4	-20%
Oct-2010	17	1	6%
Oct-2011	17	0	0%
Oct-2012	16	-1	-6%
Oct-2013	15	-1	-6%
Oct-2014	13	-2	-13%

Years shaded were not surveyed. The numbers were averaged from previous and following surveys.

The table above shows the percentage increase/decrease of pairs at Manawahe from the start of regular census operations. The years 2006 and 2008 were not surveyed and the numbers (shaded) have been averaged from the previous and following census surveys. As can be seen, the highest annual increase was 40% from 2002 to 2003; the overall annual average increase has been 7%. Unfortunately since the peak number in 2007 of 24 pair there has been an average drop of 5% per year.



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Recommendations

The following recommendations have been stated in the last three census reports. The first recommendation is the only one that has not been acted upon.

1. I recommend that if there is no significant increase in kokako numbers after the next census, intensive monitoring using video surveillance or programmed nest inspections be carried out over the nesting period. These are the most effective ways to find out what is happening during the *vulnerable* breeding season.
2. That independent specialist advice is sought from Department of Conservation and independent specialists.
3. The next population census is carried out from mid-September 2015.

References

1. Bancroft, B.; Bancroft, N. 2014: Manawahe Kokako Survey Report October 2013. *Unpublished report prepared for the Bay of Plenty Regional Council and the Manawahe Kokako Trust.*



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

2. Bancroft, B.; Bancroft, N. 2013: Manawahe Kokako Survey Report October 2012. *Unpublished report prepared for the Bay of Plenty Regional Council and the Manawahe Kokako Trust.*
3. Bancroft, B.; Bancroft, N. 2012: Manawahe Kokako Survey Report October 2011. *Unpublished report prepared for the Bay of Plenty Regional Council and the Manawahe Kokako Trust.*
4. Bancroft, B.; Bancroft, N. 2011: Manawahe Kokako Survey Report October 2010. *Unpublished report prepared for Environment Bay of Plenty and the Manawahe Kokako Trust.*
5. Bancroft, B.; Jones, G. 2007: Manawahe Kokako Survey Report October 2007. *Unpublished report prepared for Environment Bay of Plenty and the Manawahe Kokako Trust.*
6. Bancroft, B.; Te Kurapa, A. 2010: Manawahe Kokako Survey Report May 2009. *Unpublished report prepared for Environment Bay of Plenty and the Manawahe Kokako Trust.*
7. Basse, B.; Flux, I.; Innes, J. 2002: Recovery and maintenance of North Island kokako populations through pulsed pest control. *Science & Research Internal Report.*
8. Bradfield, P.; Innes, J.; Sinclair, A.R.E. Making endangered species safe: the case of the kokako of North Island, New Zealand. *Published on-line: 3 April 2006*
9. Calcott, A.; Delamare, G.; Flux, I.; Hudson, J.; Innes, J.; Molles, L.; Peters, D.; Waas, J. 2008 "Acoustic anchoring" and the successful translocation of North Island kokako (*Callaeas cinerea wilsoni*) to a New Zealand mainland management site within continuous forest., *Notornis, Vol. 55: 57-68*
10. Dennis, B. 1989. Allee effects: population growth, critical density and the chance of extinction. *Natural Resource Modeling 3: 481-537.*
11. Double, M.; Murphy, M. 2000: Genetic variation within and among populations of North Island kokako. *Science & Research Internal Report.*
12. Flux, I.; Bradfield, P.; Innes, J. 2006. Breeding biology of North Island kokako (*Callaeas cinerea wilsoni*) at Mapara Wildlife Management Reserve, King Country, New Zealand. *Notornis 53(2): 199-207.*
13. Jamieson, I.G. 2009: Loss of genetic diversity and inbreeding in New Zealand's threatened bird species. *Science for Conservation 293.*



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz


14. Jones, G.; Hudson, J. 2002: Manawahe Kokako Report May 2002. *Unpublished report prepared for Environment Bay of Plenty and the Manawahe Kokako Trust.*
15. Jones, G.; Hudson, J. 2003: Manawahe Kokako Report May 2002/2003. *Unpublished report prepared for Environment Bay of Plenty and the Manawahe Kokako Trust.*
16. Jones, G.; Hudson, J. 2004: Manawahe Kokako Report May 2003/2004. *Unpublished report prepared for Environment Bay of Plenty and the Manawahe Kokako Trust.*
17. Hudson, J. 2005: Manawahe Kokako Report May 2004/2005. *Unpublished report prepared for Environment Bay of Plenty and the Manawahe Kokako Trust.*



Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Graph 1. Manawahe Kokako Pair Total/Pair Trend Comparison 2014






Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Graph 2. Manawahe Kokako Pairs, Rat Tracking Data & Possum RTCI Over Time






Eco Rescue Ltd

Bruce Bancroft
0276318599

Map 1. Manawahe Kokako Area Surveyed 2013






Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Map 2. Kokako Pair Territory Comparison 2012-2013





Eco Rescue Ltd

Bruce Bancroft
0276318599
bruce@ecorescue.co.nz

Map 3. Kokako habitat around survey area

